

Precambrian Research

PRERES

Chec

journal homepage: www.elsevier.com/locate/precamres

A newly identified Precambrian terrane at the Pamir Plateau: The Archean basement and Neoproterozoic granitic intrusions

Chuan-Lin Zhang^{a,*}, Hai-Bo Zou^b, Xian-Tao Ye^a, Xiang-Yan Chen^a

^a College of Oceanography, Hohai University, Nanjing 210098, PR China

^b Department of Geosciences, Auburn University, Auburn, AL 36849-5305, USA

ARTICLE INFO

Keywords: Pamir syntax Precambrian terrane Zircon U-Pb ages Geochemistry Tectonic implications

ABSTRACT

A number of Precambrian terranes are distributed in the Tibetan Plateau, including the Lhas Qiangtang, Qaidam, Qilian and Tianshuihai terranes. These terranes exhibit distinct features in rock and tectonothermal history. They are important for understanding the evolution of the Tethys Oc assemblage and breakup of the Precambrian supercontinents. In this contribution, we identify a terrane at the southeastern section of the Pamir Syntax and we name it as Mazar terrane. Systematic dating shows that the basement, mainly composed of metamorphic bimodal volcanic rocks and g deposited at ca.2.5 Ga and was overprinted by ca.2.0 Ga amphibolite-facies metamorphism. The Ne granitic intrusions into the basement, emplaced at 840-835 Ma. Geochemistry of the granitic intrutwo distinct types of granitic rocks, including the trondhjemite-tonalites and granodiorites. The s jemite- tonalites show TTG-like or adakitic signatures such as enrichment of LREE and LILEs, deple and HFSE, resulting in their high Sr/Y and (La/Yb)_N ratios. The granodiorites exhibit typical sign calc-alkaline I-type granites. Both trondhjemite-tonalites and granodiorites have coupled unradio rock Nd and zircon Hf isotope compositions. ϵ Nd(T) and ϵ Hf(T) values of the trondhjemite- tonalite -9.5 to -8.7 and from -14 to -10, respectively. The granodiorites have slightly more radio isotope compositions with ϵ Nd(T) values ranging from -5.5 to -4.4 and ϵ Hf(T) values ranging is -7. According to elemental and isotope signatures of the granitic intrusions, the trondhjemite-to derived from partial melting of a mafic crust at depth \geq 30 km with a garnet amphibolite residue granodiorite was formed by partial melting of a mafic crust at a shallower level and mantle-der could be involved in the genesis of the granodiorite. In line with their geochemistry and the granodiorite is slightly younger than that of the trondhjemite-tonalites, we propose that those grani were most likely formed in an extensional environment, and that they could be part of the ear activity related to the initial breakup of the Rodinia supercontinent. Our study, in combination w works, suggests that the Mazar terrane, as well as the Qilian, Qaidam, Tianshuihai terranes, is continental fragments drifted from the Yangtze Block during the breakup of the Rodinia supercon

1. Introduction

The Tibetan Plateau is a geological amalgamation of several continental collision events since the Early Paleozoic (e.g., Dewey et al., 1988; Kapp et al., 2007; Tapponnier et al., 1981; Yin and Harrison, 2000; Yin et al., 2010a, 2010b; Zhu et al., 2012; Gibbons et al., 2015). There are several major terranes having Precambrian basement in this huge orogenic belt, such as the Lhasa, Sibumasu, Qiangtang, Qaidam, Qilian and Tianshuihai terranes (Chen et al., 2009, 2012, 2013; Zhang et al., 2010, 2012, 2013a, 2014a,b; Tung et al., 2012; Yu et al., 2017) (Fig. 1). These terranes were believed to be drifted from the Northern margin of the Great India or Australia and were gradually the Eurasia continent during the long evolution history of Ocean (Yin and Harrison, 2000). Among them, the most stu terrane recorded intensive Pan-Africa tectonothermal eve most features similar to the Pan-Africa north India orogen (2

2010, 2012; Lin et al., 2013). As for other terranes, some researchers suggested that some of them, such as Qaidam and Qilian terranes, were drifted from the Tarim Block or even the Yangtze Block (Lu et al., 2006, 2008; Song et al., 2012; Tung et al., 2013). Thus, their tectonic affinities are still ambiguous. Studies in recent years have revealed that these terranes show features distinct from each other, such as the rock

E-mail address: zchuanlin1968@gmail.com (C.-L. Zhang).

https://doi.org/10.1016/j.precamres.2017.11.006 Received 12 September 2017; Received in revised form 5 November 2017; Accepted 15 November 2017 Available online 16 November 2017 0301-9268/ © 2017 Elsevier B.V. All rights reserved.

^{*} Corresponding author.

assemblages and tectonothermal history. Thus, the detailed structure, phases of igneous activity and metamorphism of the individual terrane could shed lights on understanding the evolution of the Tethys Ocean and the evolution of global Precambrian supercontinent.

As an important part of the Tibetan Plateau, the Pamir Syntax at the northwestern end of the India-Asia collision zone underwent long-time drifting, accretion and collision since early Paleozoic (or Sinian) and the final suturing and basin closure occurred during the Mesozoic (e.g. Schwab et al., 2004; Xiao et al., 2002, 2005; Zanchi and Gaetani, 2011; Robinson et al., 2016; Robinson 2015; Angiolini et al., 2013, 2015; Rutte et al., 2017; Li et al., 2017). The most important tectonic unit in Pamir Syntax is the Tashkuergan terrane (also termed as Tashkuergan-Tianshuihai terrane), in which the amphibolite-granulite facies metamorphic rocks were identified as the Precambrian basement (Xinjiang, 1993; Yang et al., 2010). However, recent works demonstrated that the basement of the Tashkuergan terrane was early Paleozoic volcanic-sedimentary sequence, which is an accretionary wedge formed during the southward subduction of the Proto-Tethys ocean (Zhang et al., 2017a). Although some metamorphic rocks were identified as Precambrian basement of the sabsissemIcthe ofTi08. ogTi08ere (the

ando4er, au,meta-

P9ionl.,bascnr, thee0the et



Fig. 2. Geological map of the Mazar terrane showing the Archaean basement and the Neoproterozoic granitic intrusions (geochronological sample locations are marked).

types of intrusions, including gabbros, diabase, tonalite and granodiorite (Fig. 3c and d). At one outcrop, we noticed a pegmatite intruding into the gneissic mafic intrusions (Fig. 3e). Bian et al. (2013) reported two zircon U-Pb ages of the granodiorites at 855 \pm 14 Ma and 836 \pm 12 Ma. As for the mafic intrusions, no ages have been reported.

Most of the granite plutons and the mafic intrusions underwent variable degrees of deformations, leading to their gneissic structure and massive structure in the centre part of the plutons (Fig. 3c and e).

3. Petrography

3.1. The metamorphic bimodal volcanic rocks

The metamorphic bimodal volcanic rocks are composed of interbedded amphibolite and leptite of variable thickness (Fig. 3a). The leptite is grayish in color and of porphyritic texture. The phenocryst (10%) is subhedral perthite (Fig. 3f). The matrix re-crystallized and mainly comprises of fine-grained feldspar and quartz with minor biotite (Fig. 3f). Due to ductile deformation, some perthite phenocrysts show eyeball forms. The amphibolite is of gneissic structure and granularlamellar blastic texture. Rock forming minerals include hornblende (25–35%), plagioclase (45–50%), Ti-Fe oxide (10–20%), biotite (2–5%) and minor quartz (less than 2%, Fig. 3 g). According to the rock-forming minerals and the rock association observed in the field, the bimodal volcanic rock series underwent amphibolite facies metamorphism. Two leptite (*meta*-rhyolite) samples (2015D038, 2015D056) were collected for geochronological analysis (longitude-latitude locations of the samples are presented in Supplementary Table 1)

3.2. The trondhjemite-tonalites

In the field, all the granitic plutons could be defined as granodiorite according to their mineral compositions (e.g., low content of quartz). Due to their distinct geochemical features (see following discussions), we divide them into two types. The trondhjemite-tonalites consist of five plutons (Fig. 2) with a total outcropping area of $\sim 100 \text{ km}^2$. Most

of them are of gneissic structure. They are composed of 45–55% plagioclases (An = 35–45), 5–15% microcline, 15–20% quartz, 10–15% biotite, 5–10% hornblende and accessory minerals, such as zircon, apatite and ilmenite. In thin sections, most rocks show variable alterations, such as the replacement of plagioclase by sericite and replacement of biotite by epidote (Fig. 3 h). Most biotites were likely alteration products of hornblende because these biotites show pseudomorph after hornblende. Some hornblende occurs in the center of the biotite (Fig. 3 h). Fine-grained felsic aggregates and the variable-degreed foliation structure indicate the dynamic metamorphism after their emplacement (Fig. 4). Four geochronological samples were collected from four plutons (2015D029, 2015D035, 2015D037, 2015D070).

3.3. The granodiorite

Only one granodiorite pluton occurs in the western side of Mazar terrane (Fig. 2) with an outcropping area of $\sim 8 \text{ km}^2$. It intrudes into the gneissic gabbros and the basement sequence. At its fringe, chrysolite-bearing quartz-feldspar pegmatite dykes can be seen. At one outcrop, the pegmatite dyke sharply cuts the gneissic gabbro (Fig. 3e). The granodiorite mainly comprises of plagioclase (35–45%), microcline (10–20%), quartz (15–25%), K-feldspar (5–10%), biotite (5–10%), hornblende (2–5%) and accessory minerals, such as zircon, apatite and ilmenite. Similar with the scenario of the tonalites, variable degrees of alteration and foliation are commonly seen in thin sections (Fig. 3i). One granodiroite sample (2015D032-1) and one pegmatite sample (2015D032-2) were collected for age determination (Fig. 2).

4. Analytical procedures

Zircon separation was carried out using conventional magnetic and density techniques to concentrate non-magnetic, heavy fractions. Zircon grains were then hand-picked under a binocular microscope. Zircon grains were mounted in epoxy mounts which were then polished to section the crystals in half for analysis. All zircons were documented

with transmitted and reflected light micrographs as well as cathodoluminescence (CL) images to reveal their morphological appearance and internal structures. Zircon U-Pb ages and Hf isotope compositions were analyzed using the MC-LA-ICP-MS method at the Tianjin Institute of Geology and Mineral Resources (TJGMR), Chinese Geological Survey. A Neptune MC-ICP-MS coupled with a 193 nm excimer laser ablation system were used to determine zircon U-Pb ages. The laser beam diameter was 35 µm and it was operated with a frequency of 10 Hz. Every set of five sample analyses was followed by analysis of the zircon standards 91500 and eight sample analyses followed by the zircon standard GJ-1 (Jackson et al., 2004), and the glass standard NIST610 (see Hou et al., 2009). Each analysis consisted of ca. 5 s of background data acquisition and 45 s of sample data acquisition. ²⁰⁷Pb/²⁰⁶Pb, ²⁰⁶Pb/²³⁸U, ²⁰⁷Pb/²³⁵U, and ²⁰⁸Pb/²³²Th ratios were corrected for laser and instrumentally induced elemental and isotopic fractionation using zircon GJ-1 as an external standard. Common Pb was corrected using the method proposed by Andersen (2002). The U-Pb concordia plots were processed with ISOPLOT 3.0 and data are presented with 1σ errors and 95% confidence limits (Ludwig, 2003). The zircon U-Pb age data are listed in Supplementary Table 1.

Hf isotope analyses were carried out using a New Wave-193 nm ArFexcimer laser-ablation system linked to a Neptune multiple-collector inductively coupled plasma mass spectrometer (LA-MC-ICP-MS). Instrumental parameters and data acquisition followed that described by Wu et al. (2006) and Geng et al. (2011). The analyses were conducted with a beam diameter of 50 μ m, 8 Hz repetition rate with a laser power of 15 J/cm



Fig. 4. Representative cathodoluminescene images of the zircons from the Archaean *meta*-rhyolite and the Neoproterozoic granitic intrusions (see details in the text). The scale bar is $100 \,\mu$ m.

2-5%. The analytical results are presented in Supplementary Table 3.

Sr-Nd isotopes were determined using a Micromass Isoprobe Multicollector ICPMS (MC-ICPMS) at TJGMR, following the procedure described by Li et al. (2004). Measured ⁸⁷Sr/⁸⁶Sr and ¹⁴³Nd/¹⁴⁴Nd ratios were normalized to ⁸⁶Sr/⁸⁸Sr = 0.1194 and ¹⁴⁶Nd/¹⁴⁴Nd = 0.7219, respectively. The reported ⁸⁷Sr/⁸⁶Sr and ¹⁴³Nd/¹⁴⁴Nd ratios were respectively adjusted to the NBS SRM 987 standard ⁸⁷Sr/⁸⁶Sr = 0.71025 and the Shin Etsu JNdi-1 standard ¹⁴³Nd/¹⁴⁴Nd = 0.512115. Sr-Nd isotope results are listed in Table 4.

5. Analytical results

5.1. Zircon U-Pb ages

5.1.1. Age of the Archean volcanic-sedimentary sequence

Zircons from sample 2015D56 range from $80\,\mu m$ to $150\,\mu m$ in length and have length to width ratios of 1-2. Most zircon crystals are slight pinkish in color and euhedral or subhedral in form. In CL images, most of them show zoning structure. Some zircons display round form and homogenous inner structure (Fig. 4). Thirty analyses were carried out on 30 grains. Among them, six spots yield significantly older ²⁰⁶Pb/²³⁸U ages ranging from 2600 Ma to 3270 Ma and they are likely to be xenocrystals. Although four spots on the zircons of homogeneous inner texture yield discordant ²⁰⁷Pb/²⁰⁶Pb ages, their ²⁰⁶Pb/²³⁸U ages are consistent to each other within analytical errors, ranging from 1930 Ma to 1998 Ma with a mean age of 1970 \pm 41 Ma (MSWD = 7.3) (Fig. 5a). These four spots have Th contents ranging from 86 ppm to 882 ppm and U from 163 ppm to 1380 ppm with Th/U ratios from 0.5 to 1.1. According to their Th/U ratios, these four zircons were likely igneous zircons. However, their homogeneous inner structure, round form and the large range of Th and U contents share features similar to those of the ca.1.8 Ga metamorphic zircons from the Paleoproterozoic rocks at the northern margin of the Tarim Block (Th/U = 0.7-2.0, Zhang et al., 2012; Long et al., 2010; Ge et al., 2013). Previous studies have demonstrated that the Th/U ratios of metamorphic zircons are controlled by multiple factors, including the crystal growth speed, presence of fluids (Hoskin and Schaltegger, 2001). Zircons which crystallized rapidly in a relatively short time might possess high Th/U ratios up to 0.7 and even up to 3.0 (Vavra et al., 1999; Whitehouse and Kamber, 2002; Santosh et al., 2007). Collectively, we suggest that the 1970 \pm 41 Ma age represents a metamorphic event. The other twenty analyses define a good discordant line with an upper intercept age of 2523 \pm 23 Ma (MSWD = 0.5, Fig. 5b) and a meaningless lower intercept age of 1484 \pm 170 Ma, in spite of variable extents of lead loss.

Twenty-nine analyses were conducted on zircons from sample 2015D038. Among them, three spots yield concordant ages with a mean $^{206}\text{Pb}/^{238}\text{U}$ age of 2607 $\,\pm\,$ 31 Ma, likely to be xenocrystals. Four spots on the zircons of homogenous inner texture yield concordant and consistent ²⁰⁷Pb/²⁰⁶Pb and ²⁰⁶Pb/²³⁸U ages with a weighted mean $^{207}\text{Pb}/^{206}\text{Pb}$ age of 1999 \pm 21 Ma (Fig. 5c). The other 22 analyses define a discordia line with an upper intercept age of 2520 \pm 31 Ma (MSWD = 0.8, Fig. 5c). On the whole, the results of these two samples are consistent to each other. According to the inner texture and Th/U ratios (0.5–1.0), we conclude that the metamorphic bimodal volcanic sequence formed during the Neoarchean. We noticed that the age data reported by Ji et al. (2011) exhibit variable extents of radiogenic lead loss. We use their original data reported by Ji et al. (2005) and constructed a discordia line with an upper intercept age of 2521 \pm 24 Ma (MSWD = 2.1, N = 11). This age is highly consistent with the ages obtained in this study.

5.1.2. Age of the trondhjemite and tonalite

Zircons from sample 2015D029 are of short columnar forms, transparent and colorless or slight yellowish, ranging from 50 μ m to 120 μ m in length with length/width ratios of 1.5–2. Oscillatory zoning or blurry zoning were observed in CL images (Fig. 4). Thirty-two analyses were conducted on 32 zircon grains. Two spots (2015D029.8, 2015D029.26) yield older but discordant ages, likely to be xenocrystals. The other thirty analyses yield consistent results within analytical error and a weighted mean age of 840.0 \pm 3.2 Ma (Fig. 6. N = 30,



Fig. 5. Concordia of the zircon U-Pb ages of the *meta*-rhyolite from the Archaean basement of the Mazar terrane (see details in the text).

MSWD = 0.16).

Zircons from the other 3 trondhjemite-tonalite samples (2015D035, 2015D037, 2015D070) share many common features. They are translucent and colorless. Most grains are prismatic with lengths ranging from 100 to 150 μ m and length-width ratios ranging from 2:1 to 3:1 (Fig. 4). In CL images, all zircon grains display oscillatory zoning, indicating their igneous origin. Thirty-two analyses were conducted on each sample (Supplementary table 1). For sample 2015D035, one of the thirty-two analyses (spot 2015D035.7) yields Paleoproterozoic age (1902 ± 22 Ma), likely to be xenocrystal. The other thirty-one analyses show variable U (79–528 ppm) and Th (99–385 ppm) contents with Th/U ratios between 0.6 and 1.0. Their ²⁰⁶Pb/²³⁸U age and ²⁰⁷Pb/²³⁵U ages are well concordant within analytical error. They form a tight cluster in

the Concordia curve with a weighted mean 206 Pb/ 238 U age of 839.0 ± 3.1 Ma (Fig. 6b. N = 31, MSWD = 0.32). Among the thirtytwo analyses conducted on sample 2015D037, one spot (2015D037.6) yields a slightly younger age for unknown reasons (206 Pb/ 238 U age = 787 ± 9 Ma), and other 31 analyses have variable Th and U contents with high Th/U ratios ranging from 0.6 to 1.0, and yield concordant 206 Pb/ 238 U ages within analytical errors with a weighted mean age of 838.4 ± 3.1 Ma (Fig. 6c. N = 31, MSWD = 0.35). For sample 2015D070, one spot (2015D070.18) of the thirty-two analyses has Neoarchean 207 Pb/ 206 Pb age (2515 ± 20 Ma), most likely a xenocrystal from the Neoarchean basement of the Mazar terrane (Ji et al., 2011). The other thirty-one analyses yield concordant 206 Pb/ 238 U ages within analytical errors with a weighted mean age of 835.4 ± 3.1 Ma (Fig. 6d. N = 31, MSWD = 0.53).

5.1.3. Ages of the granodiorite and the pegmatite

Zircons from the granodiorite (2015D032-1) and the pegmatite (2015D032-2) at the margin of the granodiorite pluton resemble each other. They are transparent, colorless, and prismatic with lengths ranging from 100 to 200 µm and aspect ratios ranging from 2:1 to 3:1 (Fig. 4). In CL images, all zircon grains display significant oscillatory zoning and lack core-mantle structure, indicating their igneous origin (Fig. 4). One spot (spot 2015D032-1.33) was excluded during age calculation due to its younger 206 Pb/ 238 U age (796 ± 8 Ma) possibly caused by radiogenic lead loss. The other thirty-eight analyses yield concordant ²⁰⁶Pb/²³⁸U ages within analytical errors with a weighted mean age of 834.5 \pm 2.7 Ma (Fig. 6e, N = 38, MSWD = 0.36). As for the pegmatite sample (2015D032-2), one spot (2015D032-2.31) yields Mesoproterozoic ${}^{206}\text{Pb}/{}^{238}\text{U}$ age (1388 ± 15 Ma), likely to be xenocrystal. Two spots (2015D032-2.8, 2015D032-2.30) show relatively younger ²⁰⁶Pb/²³⁸U ages possibly due to variable radiogenic lead loss. The other thirty-two analyses have concordant ²⁰⁶Pb/²³⁸U ages within analytical errors with a weighted mean age of $836.2 \pm 3.3 \,\text{Ma}$ (Fig. 6 f. N = 38, MSWD = 0.21). This age is interpreted to the crystallization age of the pegmatite dyke.

5.2. Elemental geochemistry

5.2.1. The trondhjemite-tonalite

The trondhjemite-tonalite samples are intermediate to silicic in compositions with $SiO_2 = 60.6-70.3\%$ (mostly between 65% and 68%, Supplementary Table 3), metaluminous to weak peraluminous in nature with A/CNK values ranging from 0.72 to 1.09 (Fig. 7). They show a calc-alkaline trend according to the bi-variation between SiO₂ and total alkali contents (Cox et al., 1979). Their total alkali Na₂O + K₂O are between 6.20% and 8.57% and Na₂O is generally higher than K₂O with $Na_2O/K_2O = 1.5-4.3$. By CIPW calculation, in the An-Ab-Or classification diagram, all the samples are plotted into the trondhjemite and tonalite fields, which is consistent with thin section observations (Fig. 8). On the Harker diagram, Al₂O₃, TiO₂, P₂O₅, Fe₂O₃^T, CaO and MgO decrease with increasing SiO₂ (Fig. 9), whereas K₂O is not correlated with SiO₂. Na₂O decreases with SiO₂ when $65\% < SiO_2 < 68\%$, but increases with SiO_2 when $SiO_2 > 68\%$ or < 65% (Fig. 9). Their FeO^T/MgO molar ratios range from 1.5 to 2.8, indicating a low -Fe nature (Fig. 8). These rocks are relatively low in Cr (10-53 ppm) and Ni (3-21 ppm), and moderate in Yb contents (0.8-1.2 ppm) and Mg[#] values (40-52). Trace element spider diagrams and REE patterns are illustrated in Fig. 10a and b (both chondrite and Primitive mantle values were from Sun and McDonough, 1989). All samples are enriched in LILEs, such as Rb, Ba, Sr, and exhibit negative anomalies in Nb, Ta, P and Ti related to neighboring elements (Fig. 10b). They have total REE contents ranging from 104 ppm to 153 ppm and are enriched in LREE and depleted in HREE, with (La/ Yb)_N ranging from 17 to 32 (mostly between 22 and 25) and insignificant Eu anomalies (Eu*/Eu = 0.90-1.05) (Fig. 10a). Their high Sr/Y (mostly between 27 and 60) and $(La/Yb)_N$ ratios but low Y and Yb



Fig. 6. Concordia of the zircon U-Pb data of the Neoproterozoic granitic intrusions in the Mazar terrane (see details in the text).

contents are generally similar to those of modern adakites and the Archean TTG rocks (Defant and Drummond, 1990; Castllo, 2012).

5.2.2. The granodiorite

Compared with those of the trondhjemite-tonalites, the granodiorite samples have relatively lower and more restricted SiO_2 contents (63–65%). They are metaluminous in nature with A/CNK values ranging from 0.95 to 1.0 (Fig. 7). Their total alkali $Na_2O + K_2O$ ranges from 7.50% to 8.20% and their Na_2O content is similar to K_2O content. By CIPW calculation, in the An–Ab–Or classification diagram, all the samples are plotted into the granodiorite field, which is consistent with thin section observations (Fig. 8). On the Harker diagram, due to restricted chemical compositions, correlations between SiO_2 with other oxides are not significant except that Al_2O_3 and $Fe_2O_3^T$ broadly decrease with increasing SiO_2 content (Fig. 9). Their FeO^T/MgO molar ratios define their medium-Fe nature (Fig. 9) and they have significantly lower Cr (less than 3 ppm) and Ni (5–13 ppm) contents but relatively higher Yb contents (1.8–2.2 ppm), compared with those of

the trondhjemite-tonalites. As for trace elements, all samples are enriched in LILE, such as Rb, Ba and Sr, and exhibit negative anomalies in Nb, Ta, P and Ti (Fig. 10d). They have total REE contents ranging from 109 ppm to 157 ppm and are enriched in LREE and depleted in HREE, with consistent (La/Yb)_N ranging from 7 to 11 and insignificant Eu anomalies (Eu*/Eu = 0.94–1.08) (Fig. 10c). Unlike the trondhjemite-tonalites, the granodiorites have very low Sr/Y ratios (13–28) but higher HREE contents (Supplementary Table 2).

5.3. Zircon Hf isotope compositions

Zircons from the 3 trondhjemite-tonalite samples (2015D029, 2015D035, 2015D037) share similar Hf isotope compositions (Supplementary Table 2, Fig. 11a). Their ε Hf(t) values are negative, ranging from -14.1 to -9.0, and form a broad Gaussian distribution for these 3 samples (Fig. 11). Furthermore, most analyses yield 1.8–1.9 Ga $T_{DM}^{\rm Hf}$ ages and 3.0–3.2 Ga $T_{DM}^{\rm C}$ ages. On the other hand, the granodiorite sample (2015D032-1) and the pegmatite sample



Fig. 7. ACNK vs. ANK diagram showing the metaluminous signature of the trondhjemitetonalites and the granodiorite.



Fig. 8. An-Ab-Or classification diagram of the Neoproterozoic granitic intrusions in Mazar terrane.

(2015D032-2) have slightly higher zircon Hf isotope compositions compared with those of the trondhjemite-tonalites, with ϵ Hf(t) values of from -10.9 to -4.5 and from -10 to -2.9, respectively. Nevertheless, these two samples also have similar zircon Hf isotope compositions, e.g., most analyses yield 1.5–1.8 Ga T_{DM}^{Hf} ages and Neoarchean T_{DM}^{C} ages. We combine these two samples and all the analyses form a broad Gaussian distribution with a peak at ~ -6 (Fig. 11b). Both the trondhjemite-tonalite and the granodiorite have zircon Hf isotope compositions overlapping that of the Neoproterozoic detrital zircons from the southern margin of the Tarim Block (Fig. 12, Zhang et al., 2016).

5.4. Whole-rock Nd-Sr isotope compositions

Sr-Nd isotope compositions as well as the calculated (87 Sr/ 86 Sr)_i, εNd(T) and Nd model ages T_{DM} and T_{DM2} for the Neoproterozoic granitic intrusions in Mazar terrane, using the same formulation of Li et al. (2004), are presented in Supplementary Table 4. The trondhjemite-tonalite samples have a narrow range of 147 Sm/ 144 Nd (0.0930–0.1010) and 143 Nd/ 144 Nd (0.511834–0.511954) ratios, corresponding to εNd(T) values (T = 840 Ma) ranging from -9.7 to -8.2. The granodiorites have slightly more radiogenic Nd isotope compositions compared with those of the trondhjemite-tonalites (Supplementary Table 4) with εNd(T) values (T = 835 Ma) of ~ -5 . Due to variable degrees of alterations, both the trondhjemite-tonalites and granodiorites have large range of $({}^{87}\text{Sr}/{}^{86}\text{Sr})_i$ values and thus are not used for petrogenetic studies.

6. Discussions

6.1. Petrogenesis of the trondhjemite-tonalites

Chemical compositions of the trondhjemite-tonalites indicate that these rocks are not S-type, M-type, or A-type granites. Their low ferroan and calc-alkalie affinities share most characteristics of the I-type granites (Fig. 9, Frost et al., 2001). On the other hand, their high Sr/Y and (La/Yb)_N ratios, in combination with their very low HREE and HFSE contents, share most features with those of adakitic rocks (Fig. 13) (Defant et al., 2002; Wang et al., 2006), or "High Ba-Sr granites" (Tarney and Jones, 1994; Fowler and Henney, 1996; Fowler et al., 2001; Qian et al., 2002; Chen et al., 2004). Different petrogenetic mechanisms have been proposed for the generation of the adakitic granites or high Ba-Sr granites, including partial melting of a subducting oceanic slab or oceanic plateau (Defant and Drummond, 1990; Kay and Kay, 1993; Stern and Kilian, 1996; Martin et al., 2005); crustal assimilation and fractional crystallization (AFC) processes from parental basaltic magmas (Castillo, 2012); partial melting of a stalled (or dead) slab (or ocean plateau) in the mantle (Pe-Piper and Piper, 1994; Defant et al., 2002; Mungall, 2002; Qu et al., 2004); partial melting of delaminated lower crust (Kay and Kay, 1993; Xu et al., 2002; Wang et al., 2004, 2006) and partial melting of mafic rocks in the lower part of a thickened crust (Atherton and Petford, 1993; Muir et al., 1995; Petford et al., 1996; Xu et al., 2002; Zhang et al., 2007a).

Due to unradiogenic whole-rock Nd and zircon Hf isotope compositions (Supplementary Tables 3 and 4), the trondhjemite-tonalites were unlikely to have been produced by partial melting of subducted oceanic crust. This is consistent with the fact that the trondhiemite-tonalites have remarkably lower Mg[#] number and Cr and Ni contents than those of the subducted slab-derived adakites (Fig. 9). AFC processes from parental basaltic magmas can be excluded according to their significant enriched Nd-Hf isotope compositions as well as the AFC curve of mantle-derived magma (Fig. 14, Stern and Kilian, 1996). Partial melting of a stalled (or dead) slab (or ocean plateau) in the mantle, as well as partial melting of delaminated lower crust models could be also ruled out because, the resultant melt rises through a zone of mantle peridotite en route to its emplacement in the upper crust, during which the significant chemical interaction between the mantle peridotite and the crustal melt would enhance the MgO, Cr, Ni contents of the magma and change the whole-rock Nd and zircon Hf isotope compositions. The elemental and isotopic features of the trondhjemite-tonalites, such as their low MgO, Cr and Ni contents and the coupled significant unradiogenic Nd-Hf isotope compositions, are inconsistent with adakitic rocks generated in the above petrogenesis models (Fig. 9).

Numerous studies (Rapp and Watson, 1995; Rapp et al., 1999; Tate and Johnson, 2000; Petford et al., 1996; Petford and Gallagher, 2001; Wang et al., 2006) have demonstrated that mafic crustal rocks may partially melt to generate liquids with high Sr/Y and (La/Yb)_N ratios at sufficient depth (\geq 30–40 km, *i.e.* \geq 1–1.2 GPa) where garnet is stable within the residual assemblage (e.g., residues of garnet-amphibolite, amphibole-bearing eclogite and/or eclogite). This model is consistent with the low Mg[#], Cr, Ni contents and their coupled unradiogenic Nd-Hf isotope compositions of the studied rocks. In this scenario, the melting temperature can be constrained by the following two factors: (1) it should be higher than 850 °C because dehydration melting of water-unsaturated basaltic rocks could not take place when the temperature is lower than 850 °C (Rapp and Watson, 1995); (2) generally the temperature of the partial melting of the lower crust induced by the underplating of mafic magma is less than 950 °C (Tsuchiya and Kanisawa, 1994). Thus, the trondhjemite-tonalites were most likely generated by partial melting of the mafic lower crust at T = 850-950 °C



Fig. 9. Harker diagrams of the Neoproterozoic granitic intrusions in Mazar terrane (three samples reported by (Bian et al., 2013) was also included, HAS-high SiO₂ adakite, LSA-low SiO₂ adakite, the ranges of the HSA and LSA are from (Martin et al., 2005) and the ranges of the variable types of adakitic rocks are from Wang et al., 2006. See more details in the text).

and at a pressure with a residue of garnet amphibolite (Fig. 12), corresponding to a depth of 30–40 km.

6.2. Petrogenesis of the granodiorite

The I-type geochemical features of the granodiorite preclude that its primitive magma was derived from partial melting of a meta-greywacke or clay-rich precursors. Chappell and White (1974) coined the "I-type" term with the implication that the sources were igneous rocks. From experimental and theoretical perspectives, the best candidates for I-type protoliths are not mafic igneous rocks but arc volcanic rocks of intermediate composition (dacites to andesites), and possibly some relatively mafic granodiorites and tonalites that are rich in biotite and hornblende (Clemens et al., 2011). Nevertheless, due to their low SiO₂ contents and sodic nature, intermediate igneous rocks, such as tonalite and granodiorite (or intermediate metamorphic igneous rocks rich in biotite and hornblende), are not suitable protoliths because melting experiments have demonstrated the melts from these intermediate rocks are highly silicic and potassic (Patiño Douce, 1997; Li et al., 2003a,b; Clemens et al., 2011). Other competing models for the formation of calc-alkaline I-type granitic magmas include: assimilation of crustal rocks by mantle derived mafic magmas, fractionation from mafic magmas and mixing between crustal melts and mantle magmas, such as the I-type granites in Lachlan folded belt, the Central Orogenic Belt (CAOB) and South China (Wang et al., 2017). However, the hybrid model could produce significant elemental and isotope variation. Moreover, the granodiorite samples significantly deviate from the AFC trend of mantle-derived magma (Fig. 14). Thus, the hybrid model can not account for the restricted elemental compositions and the Nd-Hf isotope compositions of the granodiorite.

It has also been suggested that the I-type felsic magmas could be differentiates of large volumes of mafic magma without significant crustal contamination, even though such volumes of mafic rocks are evidently not present (Sisson et al., 2005). According to their significantly enriched whole-rock Nd and zircon Hf isotope compositions, the primitive magma of the granodiorite must be derived from hyperenriched mantle. Such hyper-enriched mantle magmas do occur (e.g., Clemens et al., 2009), but they are intermediate rather than mafic and have not fractionated to produce granites but rather represent mixing end-members (Clemens et al., 2009). Thus, this model is also not suitable for the genesis of the granodiorite although we can not rule out this mechanism completely.



Fig. 10. Chondrite-normalized REE patterns and Primitive mantle -normalized incompatible elements spider diagrams of the Neoproterozoic granitic intrusions (both the Chrondrite and Primitive mantle values are from Sun and McDonough, 1989).

We notice that the granodiorites share most major elemental and Nd-Hf isotope features with those of the trondhjemite-tonalites except for their distinct incompatible trace elements compositions (Supplementary Table 3), indicating that their possible similar magma sources but different pressure-temperature condition of partial melting. Having largely ruled out the hybrid model and mantle-derived mafic magma fractionation mechanism, the most suitable magma source for the granodiorite was Precambrian basaltic rocks. However, due to their relatively radiogenetic Nd-Hf isotope compositions and higher Mg[#] than those of the trondjemite-tonalites, mantle-derived magma could be involved in the genesis of granodiorite.

Melting experiments of Patiño Douce (1997) demonstrate that shallow dehydration melting of basaltic rocks at > 900 °C generates A-type granites, whereas deep (≥ 8 kbar) dehydration melting generates I-type granites. Nevertheless, compared with the trondhjemite-tonalites, the depth of the partial melting should be shallower than 30 km. Thus, we suggest that the granodiorites were derived from partial melting of a mafic protolith at crustal levels of pressure 8–12 kbar and T > 900 °C.

6.3. Tectonic implications

6.3.1. Tectonic environment of the Neoproterozoic granitic intrusions

It is generally accepted that calc-alkaline I-type granites are genetically related to plate subduction. However, unlike mantle-derived basaltic rocks, the geochemical features of the granites are mainly controlled by the chemical compositions of their crustal sources and pressure–temperature condition of the partial melting. Therefore, they could not be used as a highly robust indicator to decipher the tectonic settings (e.g., Li et al., 2003a). In fact, many studies in recent years have revealed that the I-type granites, even including the adakitic granites, were geodynamically related to extensional environment, such as some Neoproterozoic and Mesozoic granites in South China, the Neoproterozoic adakitic granites in northern margin of Tarim (Zhang et al., 2007a,b; Ye et al., 2008; Li et al., 2003a, 2013).

The elemental signatures, unradiogenic, homogeneous and coupled Nd-Hf isotope compositions of the trondhjemite-tonalites and the granodiorite revealed that they were derived from partial melting of distinct Precambrian basaltic rocks at different depth, excluding the possibility of mantle-derived mafic magma involvement in their genesis. Normal geothermal gradients are seldom sufficient to provide the necessary heat for partial melting of the crust. Crustal doubling, by thrusting or homogeneous thickening, also provides insufficient heat. Extra heat required to induce partial melting of the lower-middle crust is generally ascribed to mantle upwelling and lithosphere extension, possibly through the intrusion and underplating of mafic magmas (Petford and Gallagher, 2001; Clemens, 2003; Li et al., 2003a). This scenario usually takes place during the post-orogenic collapse and continental breakup (Clemens, 2003). It is noted that the granodiorite is slightly younger than the trondhjemite-tonalites. Thus, we tentatively suggest the Neoproterozoic granitic intrusions in Mazar terrane could be formed in an extensional tectonic setting. At the early stage, the mafic magma underplating induced the partial melting of the mafic crust at a depth of \geq 30 km, and produced the melts with significant adakitic or high Ba-Sr granitic elemental features similar to the voluminous 820-800 Ma adakitic rocks in northern margin of the Tarim Block (Zhang et al., 2007a,b). Due to progressive crustal extension, the partial melting at the middle crust led to the formation of the granodiorite.

6.3.2. The initial breakup of the Rodinia supercontinent

The Rodinian continents, such as Australia, Laurentia, South China, South Korea, India, Seychelles and Tarim, witnessed voluminous and diverse Neoproterozoic (ca. 830–720 Ma) continental intraplate magmatism. This global intraplate magmatism has commonly been attributed to mantle plumes or a superplume that caused the breakup and fragmentation of the supercontinent Rodinia (e.g., Heaman et al., 1992; Park et al., 1995; Li et al., 1999, 2003a, 2008a; Zhao and Cawood, 2012). Generally, initial activity of a mantle plume can produce a large



Fig. 11. Histogram of the zircon ϵ Hf(t) values of the Neoproterozoic intrusions (see details in the text).



Fig. 12. Crystallization age vs. ϵ Hf(t) diagram showing the Hf isotope compositions of the Neoproterozoic trondhjemite-tonalites and granodiorites overlapping with those of the Neoproterozoic detrital zircons in SW Tarim (Zhang et al., 2016).

quantity of basaltic magmas and form continental flood basalts (CFBs) or basaltic plateaus. The "global" onset of the Neoproterozoic mantle superplume is still a matter of debate, although most geologists suggested that the Neoproterozoic mantle plume initiated at \sim 820 Ma or a little earlier (e.g., Li et al., 1999). The \sim 850 Ma Shenwu dolerite dykes, the \sim 850 Ma Guandanshan pluton in the western Yangtze Block and the \sim 850 Ma bimodal volcanic sequence in the Cathysia Block were

Precambrian Research 304 (2018) 73-87



Fig. 13. Sr/Y vs. Y, $(La/Yb)_N$ vs. Yb_N diagrams after Defant and Drummond (1990) (see details in the text).



Fig. 14. SiO₂ vs. MgO diagram. The mantle/crustal AFC curves are after Stern and Kilian (1996) (curves 1 and 3) and Rapp et al. (1999) (curve 2). The starting points of curves 1 and 2 represent a pure slab-melt composition and metabasaltic or eclogite experimental melt (Stern and Kilian, 1996) (Rapp et al., 1999), respectively. The field of metabasaltic and eclogite melts (1–4.0 GPa) is after Stern and Kilian (1996). The fields of metabasaltic and eclogite experimental melts that are hybridized with peridotite are after Rapp et al. (1999).



Fig. 15. could J/CS03-4874i6-1.3159tf0-487likelFigH2466tz3(nb)+350.252460jn350.252460j13529101432810782(241)(3506048188.400); a 1890/8ah 239495ai, devous 1101.9(ct-35th632844213532823421353282341)(http://doi.org/10.10110/10110

interpreted as the earliest intraplate igneous activity that may be related to the Rodinia breakup (Fig. 15a) (Li et al., 2003b, 2008b, 2008c). In NW China, Song et al. (2010) reported ~850 Ma high-Ti and low-Ti series metamorphic basaltic rocks along the northern margin of the Qaidam Block, and they suggested that those eclogite-facies metamorphic basaltic rocks were part of the earliest dismembered continental flood basalt (CFB) province genetically related to the Rodinia superplume. In the southwestern Tarim Block, 850-830 Ma OIB-like basalts indicate an extensional environment (Fig. 15a, Zhang et al., 2016). 850-840 Ma igneous activities were also observed in Australia, India and South Africa (see Li et al., 2003a and references therein) and these early Neoproterozoic igneous rocks were also interpreted to represent the initial igneous rocks of the Rodinia breakup. Accordingly, we tentatively suggest that the 840-835 Ma granitic intrusions emplaced in the Mazar terrane are genetically related to the Rodinian superplume and that the Mazar terrane could be a continental fragment drifted from the Rodinian continent during its breakup.

6.3.3. Two distinct types of the Precambrian terranes in Tibetan Plateau

According to recent studies, both the Southern Kunlun terrane (SKT) in the eastern section of the Western Kunlun Orogenic Belt and the Taskuergan terrane (TAT) in NE Pamir were early Paleozoic accretionary wedges induced by the Proto-Tethys ocean subduction (Zhang et al., 2007b, 2017a, 2017b). On the other hand, the basement of the Tianshuihai terrane (TST), locally known as Tianshuihai Group, is a low-greenschist facies *meta*-graywacke deposited during the late Neoproterozoic. More importantly, detrital zircon U-Pb dating yielded three significant peaks at ca.890 Ma, ca.805 Ma and ca.750 Ma (Hu et al., 2016; Zhang et al., 2017a). Thus, the Mazar terrane between the north Pamir and the Central Pamir is the only reliable early Precambrian terrane identified in the WKOB.

The configuration of the Columbia supercontinent reconstructed by Zhao et al. (2002) and Rogers and Santosh (2002) shows that the cratonic blocks in South America and West Africa were welded by the 2.1–2.0 Transamazonia and Eburnean Orogens, the Kaapvaal and Zimbabwe Cratons in southern Africa were collided along the ~ 2.0 Ga Limpopo Belt (Rogers and Santosh, 2002, 2009). In the Yangtze Block, previous studies demonstrated that the Achaean Kongling complex was overprinted by 2.03–1.97 Ga granulite facies metamorphism (Ling et al., 2001; Zhang et al., 2006; Jiao et al., 2009). The other Paleoproterozoic orogenic events documented in Australia, North America, Atlantic and Baltica mostly took place during 1.9–1.8 Ga. The central India tectonic zone was most likely connected with the trans-North China Orogen and the collision took place at ~ 1.85 Ga, although the detailed orogenic process remains equivocal (Zhao et al., 2012).

The ca.2.0 Ga metamorphism of the basement in Mazar terrane and the ca.840 Ma granitic intrusions were also documented in Qaidam terrane and Yangtze Block (Zhang et al., 2006, 2014a,b; Chen et al., 2013). Thus, the Mazar terrane shares most features of the basement of the Yangtze Block and Qaidam terrane, but differs from those of the North China and the India.

As mentioned earlier, the tectonic affinity of the Precambrian terranes distributed in the Tibet Plateau remains foggy. Geologists unequivocally believed that the Lhasa terrane, possibly including the South Qiangtang terrane, was drifted from northern margin of the Gondwana Land due to its significant overprinting by the Pan-African metamorphism (Zhang et al., 2010, 2012; Zhu et al., 2012). As for other terranes, studies in recent years revealed that most of them, such as the Qilian and Qaidam, share most Precambrian features with those of the Tarim and Yangtze (Chen et al., 2012, 2013; Song et al., 2010; Tung et al., 2012, 2013; Zhang et al., 2013b, 2014a,b). Recent work demonstrated that the age spectra of the detrital zircons and rock association of the Tianshuihai Group are consistent with those of the late Neoproterozoic in South China (Banxi Group and its equivalences, Meng et al., 2013; Zhang et al., 2017a). The lack of the Pan-African metamorphism in these terranes indicates that they were likely located at the northern fringe of the Gondwana Land, which is in line with paleontology evidence (Metcalfe, 2011, 2013, 2017; Zhang et al., 2015; Li et al., 2017). These terranes could assemble to the fringe of the Gondwana Land during the closing of the Proto-Tethys Ocean which was formed during the breakup of the Rodinia (Li et al., 2017). Therefore, we could divide the Precambrian terranes in the Tibet into two distinct types, i.e., the Tarim-Yangtze affinity type (including the Qilian, Qaidam, Northern Qinling, Tianshuihai and Mazar) and the Gondwana affinity type (the Lhasa, South Qiangtang) (Fig. 15b).

7. Conclusions

- (1) The Mazar terrane is a newly identified Precambrian terrane in Pamir Plateau. It was mainly composed of Neoarchean metamorphic bimodal volcanic sequence and *meta*-greywackes. Neoproterozoic granitic plutons intruded the basement.
- (2) The Neoarchean metamorphic volcanic-sedimentary sequence was deposited at ca.2.5 Ga and then was overprinted by ca.2.0 Ga metamorphism, possibly related to the assemblage of the Columbia supercontinent. The trondhjemite-tonalites emplaced at ca.840 Ma and granodiorite emplaced slightly later at ca.835 Ma.
- (3) The trondhjemite-tonalites were derived from partial melting of a

mafic crust at depth \geq 30 km whereas the granodiorite formed by partial melting of a mafic crust at shallower level than that of the trondhjemite-tonalites. Their ages and geochemical features suggest that both of them likely formed at an extensional environment.

(4) The 840–835 Ma granitic intrusions distributed in Mazar terrane could be part of the earliest igneous activity related to the initial breakup of the Rodinia supercontinent. The Mazar terrane, as well as the Qilian, Qaidam and Tianshuihai terranes, was likely a continental fragment drifted from Yangtze during its breakup.

Acknowledgements

We sincerely thank Mr. Wen-Hua Ji for his help with the field work and supplying the 1/50000 geological maps. We are grateful to Hong-Ying Zhou for his assistance with LA-ICP-MS zircon dating, and to Zhang Jian for his help with zircon Lu-Hf isotope analyses. This project is funded by the National 305 Project of China (2015BAB05B0X-03) and the Fundamental Research Fund for Central Universities (B16020127). We are grateful to Dr. Wei Wang and an anonymous reviewer for their constructive reviews that greatly improved the quality of this paper.

Appendix A. Supplementary data

Precambrian Research 304 (2018) 73–87

Yangtze Block, South China. Chin. Sci. Bull. 46 (4), 326-329.

- Long, X.P., Yuan, C., Sun, M., Zhao, G.C., Xiao, W.J., Wang, Y.J., Yang, Y.H., Hu, A.Q., 2010. Archean crustal evolution of the northern Tarim Craton, NW China: Zircon U-Pb and Hf isotopic constraints. Precambr. Res. 180, 272–284.
- Ludwig KR. 2003. User's manual for Isoplot 3.00:a geochronological toolkit for Microsoft Excel. Kenneth R. Ludwig.
- Lu, S.N., Yu, H.F., Li, H.K., 2006. Research on Precambrian Major Problems in China. Geol. Pub. House 1–206 (in Chinese).
- Lu, S.N., Li, H.K., Zhang, C.L., Niu, G.H., 2008. Geological and geochronological evidence for the Precambrian evolution of the Tarim Craton and surrounding continental fragments. Precambr. Res. 160, 94–107.
- Martin, H., Smithies, R.H., Rapp, R., Moyen, J.F., Champion, D., 2005. An overview of adakite, tonalite-trondhjemite-granodiorite (TTG) and sanukitoid: relationships and some implications for crustal evolution. Lithos 79, 1–24.
- Metcalfe, I., 2011. Palaeozoic-Mesozoic History of SE Asia. In: Hall, R., Cottam, M., Wilson, M. (Eds.), The SE Asian Gateway: History and Tectonics of Australia-Asia Collision. Geological Society of London Special Publications 355, pp. 7–35.
- Metcalfe, I., 2013. Gondwana dispersion and Asian accretion: Tectonic and palaeogeographic evolution of eastern Tethys. J. Asian Earth Sci. 66, 1–13.
- Meng, Q.X., Zhang, J., Geng, J.Z., Zhang, C.L., Huang, W.C., 2013. Zircon U-Pb age and Hf isotope compositions of Lengjiaxi and Baxi Groups in middle Hunan Province: implications for the Neoproterozoic tectonic evolution in South China. Geol. China 40, 191–216 (in Chinese with English abstract).
- Muir, R.J., Weaver, S.D., Bradshaw, J.D., Eby, G.N., Evans, J.A., 1995. Geochemistry of the Cretaceous separation point batholith, New Zealand: granite magmas formed by melting of mafic lithosphere. J. Geol. Soc., Lon. 152, 689–701.
- Mungall, J.E., 2002. Roasting the mantle: slab melting and the genesis of major Au and Au-rich Cu deposits. Geology 30, 915–918.
- Nowell, G.M., Kempton, P.D., Noble, S.R., Fitton, J.G., Saunders, A.D., Mahoney, J.J., Taylor, R.N., 1998. High precision Hf isotope measurements of MORB and OIB by thermal ionisation mass spectrometry: insights into the depleted mantle. Chem. Geol. 149, 211–233.
- Park, J.K., Buchan, K.L., Harlan, S.S., 1995. A proposed giant radiating dyke swarm fragmented by the separation of Laurentia and Australia based on paleomagnetism of ca. 780 Ma mafic intrusions in western North America. Earth Planet. Sci. Lett. 132, 129–139.
- Patiño Douce, A.E., 1997. Generation of metaluminous A-type granites by low-pressure melting of calc-alkaline granitoids. Geology 25, 743–746.
- Petford, N., Gallagher, G., 2001. Partial melting of mafic (amphibolitic) lower crust by periodic influx of basaltic magma. Earth Planet. Sci. Lett. 193, 483–499.
- Petford, N., Atherton, M.P., Halliday, A.N., 1996. Rapid magma production rates, underplating and remelting in the Andes: isotopic evidence from northern-central Peru (90–110s). J. South. Am. Earth Sci. 9, 69–78.
- Pe-Piper, G., Piper, J.W., 1994. Miocene magnesium andesites and dacites, Evia, Greece: adakites associated with subducting slab detachment and extension. Lithos 31, 125–140.
- Qian, Q., Chung, S.L., Li, Y.T., Wen, D.R., 2002. Geochemical characteristics and petrogenesis of the Badaling high Ba–Sr granites: a comparison of igneous rocks from North China and Dabie-Sulu Orogen. Acta Geol. Sinica 18, 275–292 (in Chinese with English abstract).
- Qu, X.M., Hou, Z.Q., Li, Y.G., 2004. Melt components derived from a subducted slab in late orogenic ore-bearing porphyries in the Gangdese copper belt, southern Tibetau plateau. Lithos 74, 131–148.
- Qu, J.F., Zhang, L.F., Ai, Y.L., Lu, Z., Wang, J.P., Zhou, H., Wang, S.Y., 2007. Highpressure granulite from Western Kunlun, northwestern China: its metamorphic evolution, zircon SHRIMP U-Pb ages and tectonic implication. Sci. China D-ser. 50, 961–971.
- Rapp, R.P., Watson, E.B., 1995. Dehydration melting of matabasalt at 8–32 kbar: implications for continental growth and crust–mantle recycling. J. Petrol. 36, 891–931.
- Rapp, R.P., Shimizu, N., Norman, M.D., Applegate, G.S., 1999. Reaction between slabderived melts and peridotites in the mantle wedge: experimental constraints at 3.8 GPa. Chem. Geol. 160, 335–356.
- Robinson, A.C., Yin, A., Manning, C.E., Harrison, T.M., Zhang, S.-H., Wang, X.-F., 2007. Cenozoic evolution of the eastern Pamir: implications for strain accommodation mechanisms at the western end of the Himalayan-Tibetan orogen. GSA Bull. 119, 882–896.
- Robinson, A.C., Ducea, M., Lapen, T.J., 2012. Detrital zircon and isotopic constraints on the crustal architecture and tectonic evolution of the northeastern Pamir. Tectonics 31, 1–16.
- Robinson, A.C., 2015. Mesozoic tectonics of the Gondwanan terranes of the Pamir Plateau. J. Asian Earth Sci. 102, 170–179.
- Robinson, A.C., Ducea, M., Lapen, T.J., 2016. Detrital zircon and isotopic constraints on the crustal architecture and tectonic evolution of the northeastern Pamir. Tectonics 31, 1–16.
- Rogers, J.J.W., Santosh, M., 2002. Configuration of Columbia, a Mesoproterozoic supercontinent. Gondwana Res. 5, 5–22.
- Rogers, J.J.W., Santosh, M., 2009. Tectonics and surface effects of the supercontinent Columbia. Gondwana Res. 15, 373–380.
- Rutte, D., Ratschbacher, L., Schneider, S., Stübner, K., Stearns, Gulzar M.A., Hacker, B.R., 2017. Building the Pamir-Tibetan Plateau—Crustal stacking, extensional collapse, and lateral extrusion in the Central Pamir: 1. Geometry and kinematics. Tectonics 36, 1–43.
- Santosh, M., Wilde, S., Li, J.H., 2007. Timing of Paleoproterozoic ultrahightemperature metamorphism in the North China Craton: evidence from SHRIMP U-Pb zircon geochronology. Precambr. Res. 159, 178–196.

Schwab, M., Ratschbacher, L., Siebel, W., Williams, M.M., Minaev, V., Lutkov, V., Chen,

F., Stanek, K., Nelson, B., Frisch, W., Wooden, J.L., 2004. Assembly of the Pamirs: age and origin of magmatic belts from the southern Tien Shan to the southern Pamirs and their relation to Tibet. Tectonics 23, 1–31.

- Scherer, E., Munker, C., Mezger, K., 2001. Calibration of the lutetium–hafnium clock. Science 293, 683–687.
- Sisson, T.W., Ratajeski, K., Hankins, W.B., Glazner, A.F., 2005. Voluminous granitic magmas from common basaltic sources. Contrib. Miner. Petrol. 148, 635–661.
- Song, S.G., Su, L., Li, X.H., Zhang, G.B., Niu, Y.L., Zhang, L.F., 2010. Tracing the 850-Ma continental flood basalts from a piece of subducted continental crust in the North Qaidam UHPM belt, NW China. Precambr. Res. 183, 805–816.
- Song, S.G., Su, L., Li, X.H., Niu, Y.L., Zhang, L.G., 2012. Grenville-age orogenesis in the Qaidam-Qilian block: the link between SouthChina and Tarim. Precambr. Res. 220–221. 9–22.
- Stern, G.R., Kilian, R., 1996. Role of the subducted slab, mantle wedge and continental crust in the generation of adakites from the Austral Volcanic Zone. Contrib. Miner. Petrol. 123, 263–281.
- Sun, S.S., McDonough, W.F., 1989. Chemical and isotopic systematics of oceanic basalt: implication for mantle composition and processes. In: Saunders, A.D., Morry, M.J. (Eds.), Magmatism in the Ocean Basin, vol. 42. Geological Society of London Special Publication, pp. 528–548.
- Tapponnier, P., Mattauer, M., Proust, F., Cassaigneau, C., 1981. Mesozoic ophiolites, sutures, and large-sale tectonic movements in Afghanistan. Earth Planet. Sci. Lett. 52, 355–371.
- Tarney, J., Jones, C.E., 1994. Trace element geochemistry of orogenic igneous rocks and crustal growth models. J. Geol. Soc. Lon. 151, 855–868.
- Tate, M.C., Johnson, S.E., 2000. Subvolcanic and deep-crustal tonalite genesis beneath the Mexican Peninsular Ranges. J. Geol. 108, 721–728.
- Tsuchiya, N., Kanisawa, S., 1994. Early Cretaceous Sr-rich silicic magmatism by slab melting in the Kitakami Mountains, Northeastern Japan. J. Geophys. Res. 99, 22205–22220.
- Tung, G.A., Yang, H.Y., Liu, D.Y., Zhang, J.X., Yang, H.J., Shau, Y.H., Tseng, C.Y., 2012. The amphibolite-facies metamorphosed mafic rocks from the Maxianshan area, Qilian block, NW China: A record of early Neoproterozoic arc magmatism. J. Asian Earth Sci. 46, 177–189.
- Tung, G.A., Yang, H.Y., Liu, D.Y., Zhang, J.X., Yang, H.J., Shau, Y.H., Tseng, C.Y., 2013. The Neoproterozoic granitoids from the Qilian block, NW China: Evidence for a link between the Qilian and South China blocks. Precambr. Res. 235, 163–189.
- Valli, F., Leloup, P.H., Paquette, J.L., et al., 2008. New U/Pb constraints on timing of shearing and long-term slip-rate on the Karakorum fault. Tectonics 27, 1–33.
- Vavra, G., Schmid, R., Gebauer, D., 1999. Internal morphology, habit and U-Th-Pb microanalysis of amphibole to granulite facies zircon: Geochronology of the Ivren Zone (Southern Alps). Contrib. Mineral. Petrol. 134, 380–404.
- Wang, Q., Zhao, Z.H., Bao, Z.W., Xu, J.F., Liu, W., Li, C.F., 2004. Geochemistry and petrogenesis of the Tongshankou and Yinzu adaktic intrusive rocks and the associated porphyry copper molybdenum mineralization in southeast Hubei, east China. Resour. Geol. 54, 137–152.
- Wang, Q., Xu, J.F., Jian, P., Bao, Z.W., Zhao, Z.H., Li, C.F., Xiong, X.L., Ma, J.L., 2006. Petrogenesis of adaktic porphyries in an extensional tectonic setting, Dexing, south China: implications for the genesis of porphyry copper mineralization. J. Petrol. 47, 119–144.
- Wang, W., Liu, S.W., Feng, Y.G., Li, Q.G., Wu, F.H., Wang, Z.Q., Wang, R.T., Yang, P.T., 2012. Chronology, petrogenesis and tectonic setting of the Neoproterozoic Tongchang dioritic pluton at the northwestern margin of the Yangtze Block: constraints from geochemistry and zircon U-Pb–Hf isotopic systematics. Gondwana Res. 22, 699–716.
- Wang, Z.M., Han, C.M., Xiao, W.J., Sun, B.X., Ding, J.X., Song, S.H., 2017. Geochemical and zircon U-Pb and Lu-Hf isotopic constraints on the origin of supracrustal rocks from the mid-Qilian terrane: a comparison between supracrustal rocks on the two sides of the eastern segment of the Altyn Tagh Fault. Precambr. Res. 294, 284–306.
- Wu, F.Y., Yang, Y.H., Xie, L.W., Yang, J.H., Xu, P., 2006. Hf isotopic compositions of the standard zircons and baddeleyites used in U-Pb geochronology. Chem. Geol. 234, 105–126.
- Whitehouse, M.J., Kamber, B.S., 2002. On the overabundance of light rare earth elements in terrestrial zircons and its implication for Earth's earliest magmatic differentiation. Earth Planet. Sci. Lett. 204, 333–346.
- Xiao, W.J., Windley, B.F., Hao, J., Li, J.L., 2002. Arc-ophiolite obduction in the western Kunlun range (China): implications for the Palaeozoic evolution of central Asia. J. Geol. Soc. 159, 517–528.
- Xiao, W.J., Windley, B.F., Liu, D.Y., Jian, P., Liu, C.Z., Yuan, C., Sun, M., 2005. Accretionary tectonics of the western Kunlun orogen, China: A Paleozoic-Early Mesozoic, long-lived active continental margin with implications for the growth of southern Eurasia. J. Geol. 113, 687–705.
- Xinjiang, BGMR., 1993. Regional Geology of the Xinjiang Uygur Autonomous Region: Beijing, Geological Publishing House. pp. 155–231 (in Chinese).
- Xu, J.F., Shinjo, R., Defant, M.J., Wang, Q., Rapp, R.P., 2002. Origin of Mesozoic adakitic intrusive rocks in the Ningzhen area of east China: partial melting of delaminated lower continental crust? Geology 30, 1111–1114.
- Yang, W.Q., Liu, L., Cao, Y.T., Wang, C., He, S.P., Li, R.S., Zhu, X.H., 2010. Geochronological evidence of Indosinian (high-pressure) metamorphic event and its tectonic significance in Taxkorgan area of the Western Kunlun Mountains, NW China. Sci. China D-ser. 53, 1455–1459.
- Ye, H.M., Li, X.H., Li, Z.X., Zhang, C.L., 2008. Age and origin of the high Ba-Sr granitoids from northern Qinghai-Tibet Plateau: implications for the early Paleozoic tectonic evolution of the Western Kunlun orogenic belt. Gondwana Res. 13, 126–138.
- Yin, A., Harrison, T.M., 2000. Geologic evolution of the HimalayanTibetan Orogen. Annu. Rev. Earth Plan. Sci. 28, 211–280.

- Yin, A., Dubey, C.S., Webb, A.A.G., Kelty, T.K., Grove, M., Gehrels, G.E., Burgess, W.P., 2010a. Geological correlation of the Himalayan orogen and Indian craton. Part 1. Structural geology, U-Pb zircon geochronology, and tectonic evolution of theShillong Plateau and its neighbouring regions in NE India. GSA Bull. 122, 336–359.
- Yin, A., Dubey, C.S., Kelty, T.K., Webb, A.A.G., Harrison, T.M., Chou, C.Y., Celerier, J., 2010b. Geological correlation of the Himalayan orogen and Indian craton. Part 2. Structural geology, geochronology, and tectonic evolution of the eastern Himalaya. GSA Bull. 122, 360–395.
- Yu, S.Y., Zhang, J.X., Li, S.Z., Sun, D.Y., Li, Y.S., Liu, X., Guo, L.L., Suo, Y.H., Peng, Y.B., Zhao, X.L., 2017. Paleoproterozoic granulite-facies metamorphism and anatexis in the Oulongbuluke Block, NW China: Respond to assembly of the Columbia supercontinent. Precambr. Res. 291, 42–62.
- Zanchi, A., Poli, S., Fumagalli, P., Gaetani, M., 2000. Mantle exhumation along the Tirich Mir Fault Zone, NW Pakistan: Pre-mid-Cretaceous accretion of the Karakoram terrane to the Asian margin, in Tectonics of the Nanga Parbat Syntaxis and the Western Himalaya, edited by M. A. Khan et al., Geol. Soc. Spec. Publ., 170, 219–236.
- Zanchi, A., Gaetani, M., 2011. The geology of the Karakoram range, Pakistan: the new 1:100,000 geological map of Central-Western Karakoram. Italy J. Geosci. 130, 161–262.
- Zhao, G.C., Cawood, P.A., Wilde, S.A., Sun, M., 2002. Review of global 2.1–1. 8 Ga orogens: implications for a pre-Rodinia supercontinent. Earth-Sci. Rev. 59, 125–162.
- Zhao, G.C., Cawood, P.A., Wilde, S.A., Sun, M., Zhang, J., He, Y.H., Yin, C.Q., 2012. Amalgamation of the North China Craton: key issues and discussion. Precambr. Res. 222–223, 55–76.
- Zhao, G.C., Cawood, P.A., 2012. Precambrian geology of China. Precambr. Res. 222–223, 13–54.
- Zhang, S.B., Zheng, Y.F., Wu, Y.B., Zhao, Z.F., Gao, S., Wu, F.Y., 2006. Zircon U-Pb age and Hf isotope evidence for 3.8 Ga crustal remnant and episodic reworking of Archean crust in South China. Earth Planet. Sci. Lett. 252, 56–71.
- Zhang, C.L., Li, X.H., Li, Z.X., Lu, S.N., Ye, H.M., Li, H.M., 2007a. Neoproterozoic ultramafic-mafic-carbonatite complex and granitoids in Quruqtagh of northeastern Tarim Block, western China: geochronology, geochemistry and tectonic implications. Precambr. Res. 152, 149–169.
- Zhang, C.L., Lu, S.N., Yu, H.F., Ye, H.M., 2007b. Tectonic evolution of Western Orogenic belt: evidences from zircon SHRIMP and LA-ICP-MS U-Pb ages. Sci. China (D) 50, 1–12.

Zhang, C.L., Li, H.K., Zou, H.B., 2013b. Tectonic framework and evolution of the Tarim

Block in NW China. Gondwana Res. 23, 1306-1315.

- Zhang, C.L., Zou, H.B., Santosh, M., Ye, X.T., Li, H.K., 2014a. Is the Precambrian basement of the Tarim Craton in NW China composed of discrete terranes? Precambr. Res. 254, 226–244.
- Zhang, C.L., Santosh, M., Zhu, Q.B., Chen, X.Y., Huang, W.C., 2015. The Gondwana connection of South China: evidence from monazite and zircon geochronology in the Cathaysia Block. Gondwana Res. 28, 1137–1151.
- Zhang, C.L., Ye, X.T., Zou, H.B., Chen, X.Y., 2016. Neoproterozoic sedimentary basin evolution in southwestern Tarim, NW China: new evidence from field observations, detrital zircon U-Pb ages and Hf isotope compositions. Precambr. Res. 280, 31–45.
- Zhang, C.L., Zou, H.B., Ye, X.T., Chen, X.Y., 2017a. Tectonic evolution of the eastern section of Western Kunlun Orogenic Belt at the northern margin of the Tibetan Plateau: new evidence from field observations and geochronology. Gondwana Res under review.
- Zhang, C.L., Zou, H.B., Ye, X.T., Chen, X.Y., 2017b. Tectonic evolution of the NE section of the Pamir Plateau: new evidences from field observation and zircon U-Pb geochronology. Tectonophysics under review.
- Zhang, J.X., Yu, S.Y., Gong, J.H., Li, H.K., Hou, K.J., 2013a. The latest Neoarchean –Paleoproterozoic evolution of the Dunhuang block, eastern Tarim craton, northwestern China: Evidence from zircon U-Pb dating and Hf isotopicanalyses. Precambr. Res. 226, 21–42.
- Zhang, L., Wang, Q.Y., Chen, N.S., Sun, M., Santosh, M., Ba, J., 2014b. Geochemistry and detrital zircon U-Pb and Hf isotopes of the paragneiss suite from the Quanji massif, SE Tarim Craton: implications for Paleoproterozoic tectonics in NWChina. J. Asian Earth Sci. 95, 33–50.
- Zhang, Z.M., Zhao, G.C., Santosh, M., Wang, J.L., Dong, X., Liou, J.G., 2010. Two stages of granulite facies metamorphism in the eastern Himalayan syntaxis, south Tibet: petrology, zircon geochronology and implications for the subduction of Neo-Tethys and the Indian continent beneath Asia. J. Metamorph. Geol. 28, 719–733.
- Zhang, Z.M., Dong, X., Santosh, M., Liu, F., Wang, W., Yiu, F., He, Z.Y., Shen, K., 2012. Petrology and geochronology of the Namche Barwa Complex in the eastern Himalayan syntaxis, Tibet: Constraints on the origin and evolution of the northeastern margin of the Indian Craton. Gondwana Res. 21, 123–137.
- Zhu, D.C., Zhao, Z.D., Niu, Y.L., Dilek, Y., Wang, Q., Ji, W.H., Dong, G.C., Sui, Q.L., Liu, Y.S., Yuan, H.L., Mo, X.X., 2012. Cambrian bimodal volcanism in the Lhasa Terrane, southern Tibet: record of an early Paleozoic Andean-type magmatic arc in the Australian proto-Tethyan margin. Chem. Geol. 328, 290–308.