

The origin of the *c.* 1.7 Ga gabbroic intrusion in the Hekou area, SW China: constraints from SIMS U–Pb zircon geochronology and elemental and Nd isotopic geochemistry

WEI-GUANG ZHU, ZHONG-JIE BAI, HONG ZHONG, XIAN-TAO YE
& HONG-PENG FAN

State Key Laboratory of Ore Deposit Geochemistry, Institute of Geochemistry, Chinese Academy of Sciences, 99 West Lincheng Road, Guiyang 550081, China

(Received 2 June 2015; accepted 15 December 2015; first published online 9 February 2016)

Abstract – The late Palaeoproterozoic to early Mesoproterozoic igneous rocks of southwestern China are characterized by a number of mafic intrusions and dykes. However, the origin and tectonic implications of these mafic intrusions and dykes remain unclear. The Hekou mafic intrusion, intruding into the Hekou Group in the Hekou area, SW China, is the biggest and most representative one. The intrusion is mainly composed of coarse-grained in the central zone (CZ) and medium- to fine-grained gabbroic rocks in the outer zone (OZ). Cameca secondary ion mass spectroscopy (SIMS) U–Pb zircon ages, and geochemical and Nd isotopic results for the intrusion are reported in this paper. SIMS U–Pb zircon ages indicate that the gabbroic rocks from the CZ and OZ were emplaced at 1735 ± 6.5 Ma and 1736 ± 4.0 Ma, respectively. This suggests that the Hekou intrusion originated from *c.* 1.7 Ga mafic magmatism in the southwestern Yangtze Block. The coarse-grained rocks in the CZ of the intrusion show fairly homogeneous major- and trace-element compositions. In contrast, the medium- to fine-grained rocks from the OZ display slightly evolved compositions, with relatively lower Mg, Mn, MgO, Al₂O₃, Cr and Ni contents, and higher SiO₂, CaO and Zr concentrations than those of the rocks from the CZ. Although the gabbroic rocks of the intrusion have low total rare earth element (REE) contents (REE = 29.3–40.2 ppm) with slightly light REE (LREE)-enriched and heavy REE (HREE)-depleted patterns, they exhibit distinct trace-element and Nd isotopic features. The rocks from the CZ are characterized by slightly LREE-enriched and ‘convex upwards’ incompatible trace-element patterns with significant Th depletion and insignificant Nb and Ta depletion relative to La. However, the rocks from the OZ have relatively flatter REE patterns than those of the rocks from the CZ. In addition, the rocks from the OZ are slightly enriched in Th and depleted in Nb and Ta relative to La. The $\epsilon_{\text{Nd}}(T)$ values of the CZ and the OZ rocks are +0.70 to +2.3 and –0.30 to +0.24, respectively. The parental magma for the Hekou gabbroic intrusion exhibits affinity with a subalkaline basaltic magma, which was possibly generated by relatively high degrees of partial melting of a slightly depleted asthenospheric mantle source. Their geochemical and isotopic variations were due to slight crystal fractionation with varying degrees of crustal contamination. The Hekou intrusion was therefore supposed to form in an anorogenic extensional environment. It is further suggested that *c.* 1.7 Ga is an important onset timing of widespread anorogenic magmatism in the southwestern Yangtze Block. We interpret the late Palaeoproterozoic gabbroic intrusion to represent anorogenic mafic magmatism, which was most likely related to the break-up of the Columbia supercontinent.

Keywords: late Palaeoproterozoic, gabbroic intrusion, zircon geochronology, anorogenic magmatism, southwestern Yangtze Block, Columbia supercontinent.

1. Introduction

The South China Craton, including the Yangtze Block to the northwest and the Cathaysia Block to the southeast (present coordinates), was presumably the central part of the Precambrian supercontinent Rodinia (Fig. 1a) (Li, Zhang & Powell, 1995). Apart from the Kongling Complex, there have been three other major rock units previously reported from the Yangtze Block, including some volcanic and meta-volcanic rocks of late Palaeoproterozoic to Mesoproterozoic age. The *c.* 1.75–1.66 Ga Dahongshan Group (marked ‘2’ in Fig. 1a) consists of meta-volcaniclastic rocks,

meta-basalts, meta-siliciclastic rocks and marble (e.g. Greentree & Li, 2008; Zhao *et al.* 2010). The Tianli schists (marked ‘3’ in Fig. 1a) represent a clastic sedimentary succession, probably formed on the southern continental shelf of the Yangtze Block between *c.* 1530 Ma and 1042 Ma (Li *et al.* 2007). The Kunyang–Huili groups (marked ‘4’ in Fig. 1a) consist of a sequence of greenschist-facies meta-sedimentary and meta-volcanic rocks that probably developed between 1.8 Ga and 1.0 Ga (Greentree *et al.* 2006; Greentree & Li, 2008; Zhao *et al.* 2010).

The Yangtze Block played an important role in the reconstruction of the Rodinia supercontinent (e.g. Li, Zhang & Powell, 1995; Li *et al.* 2002, 2008), but little is known as to whether or not it was also a component

Author for correspondence: zhuweiguang@vip.gyig.ac.cn




Figure 1. (Colour online) (a) Simplified tectonic map showing the study area in relation to South China's major tectonic units (Li *et al.* 2007). Numbers 1–4 indicate the Kongling Complex, the Dahongshan Group, the Tianli schists and the Kunyang–Dongchuan groups, respectively; (b) geological map of the late Palaeo- to Mesoproterozoic strata and Proterozoic intrusions in the Kangdian region, SW China (modified from Wu *et al.* 1990; Zhao & Zhou, 2011).

of the Palaeo- to Mesoproterozoic supercontinent called Nuna (Hoffman, 1989, 1997; Evans & Mitchell, 2011; Zhang *et al.* 2012) or Columbia (Rogers & Santosh, 2002; Zhao *et al.* 2002a; Hou *et al.* 2008b; Meert, 2012; we will use this name in the remainder of this paper for the late Palaeo- to Mesoproterozoic supercontinent). Nonetheless, 2.03–1.97 Ga granulites and 1.85 Ga A-type granites and mafic dykes have been identified in the northern part of the Yangtze Block (the Kongling Complex; ‘1’ in Fig. 1a) (Zhang *et al.* 2006a; Sun *et al.* 2008; Wu *et al.* 2008; Xiong *et al.* 2009; Peng *et al.* 2009, 2012), which coincide with the age of the Columbia supercontinent assembly (Hoffman, 1989, 1997), and may therefore be genetically linked to the evolution of the supercontinent. Yao, Shu & Santosh (2011) presented evidence from detrital zircon U–Pb geochronology, Hf isotopes and geochemistry from the Cathaysia Block (Fig. 1a) with clear age peaks at 1930–1520 Ma, which they correlated with the Columbia supercontinent. In addition, a number of studies have documented the late Palaeoproterozoic to early Mesoproterozoic igneous rocks in the southwestern Yangtze Block in recent years. The presence of late Palaeoproterozoic to early Mesoproterozoic (1.7–1.5 Ga) mafic magmatism in the southwestern Yangtze Block is contemporaneous with the break-up of the Columbia supercontinent, and provides further evidence to support the idea that the Yangtze Block was likely a part of the Columbia supercontinent (Zhao *et al.* 2010, 2013; Fan *et al.* 2013). Meanwhile, the Yinmin Formation of the Lower Kunyang Group in the southwestern Yangtze Block contains zircons of ages dominantly from late Archaean to Palaeoproterozoic (2.8–2.7 Ga, 2.5–2.3 Ga and *c.* 1.85 Ga). The Heishantou Formation of the Upper Kunyang Group has two major age populations, *i.e.* 1.8–1.6 Ga and *c.* 1.0 Ga. The presence of abundant *c.* 1.85 Ga zircons also suggests that the Yangtze Block was probably a part of the Columbia supercontinent during Palaeoproterozoic time (Wang *et al.* 2012).

In recent years, many intrusive igneous rocks in the southwestern Yangtze Block have been gradually identified in age as late Palaeoproterozoic to early Mesoproterozoic, although these intrusive igneous rocks in the Kangdian region are sparse and on a small scale. For example, a number of mafic intrusions and dykes intruded into the Dahongshan Group, Lower Kunyang Group (also called the Dongchuan Group; we will use the name Lower Kunyang Group in the remainder of the paper), Hekou Group and Huili Group in the Kangdian region (Zhao *et al.* 2010, 2013; Guan *et al.* 2011; Zhao & Zhou, 2011; Fan *et al.* 2013). A dolerite dyke cutting the Dahongshan Group has a laser ablation inductively coupled plasma mass spectrometry (LA-ICP-MS) U–Pb zircon age of 1659 ± 16 Ma (Zhao & Zhou, 2011). The other dolerite dykes that cut the Yinmin Formation of the Lower Kunyang Group have LA-ICP-MS and a secondary ion mass spectroscopy (SIMS) U–Pb zircon ages of 1690 ± 32 Ma (Zhao *et al.* 2010) and 1701 ± 28 Ma (Zhao *et al.* 2013), respectively.

In addition, the Zhuqing Fe–Ti–V oxide ore-bearing mafic intrusions that intruded into the Huili Group were dated at 1494 ± 6 Ma (zircon U–Pb), 1486 ± 3 Ma and 1490 ± 4 Ma (baddeleyite U–Pb), respectively (Fan *et al.* 2013). However, owing to the lack of more systematic geochemical and isotopic data, the origin and tectonic implications of these mafic dykes and intrusions remain unclear. Thus, further studies on the timing and tectonic setting of these mafic dykes and intrusions are needed, which will play an important role in elucidating the late Palaeoproterozoic to early Mesoproterozoic tectonic evolution of the southwestern Yangtze Block.

There are numerous mafic intrusions that intruded the Hekou Group in the Hekou area, SW Huili County (D. F. He, unpub. Ph.D. thesis, Graduate School of the Chinese Academy of Sciences, 2009; Zhou *et al.* 2009; Guan *et al.* 2011; Chen & Zhou, 2012; Chen, Zhou & Zhao, 2013). Recent studies have focused on the ages of these intrusions. The *c.* 1.7 Ga dolerite and gabbroic intrusions that intruded into the Hekou Group were recently identified in the Hekou area (Guan *et al.* 2011; Chen, Zhou & Zhao, 2013). However, Guan *et al.* (2011) reported a SHRIMP zircon age of 1710 ± 8 Ma for a dolerite intrusion that intruded into the Hekou Group, which is very different from the whole-rock Sm–Nd isochron age of *c.* 850 Ma for some gabbroic and dolerite intrusions in this area (Zhou *et al.* 2009). In addition, the petrogenesis of the mafic intrusions is not well understood in this area. Thus, systematic geochronological, elemental and Nd isotopic investigations on these rocks are further required to elucidate the origin and tectonic implications of these mafic intrusions.

In this paper, we report precise Cameca SIMS U–Pb zircon ages, and geochemical and Nd isotopic data for the biggest and most representative Hekou gabbroic intrusion in the Hekou area with the aim to (1) date the crystallization age for the gabbroic intrusion; (2) constrain the origin and petrogenesis of the intrusion; and (3) shed new light on the late Palaeoproterozoic tectonic evolution in the southwestern Yangtze Block. Our new results confirm that the gabbroic intrusion emplaced at *c.* 1.7 Ga in the Hekou area was derived from a slightly depleted mantle source and formed in an intracontinental rift, most likely related to the break-up of the Columbia supercontinent.

2. Geological background and petrography

The South China Craton consists of two major Precambrian blocks: the Yangtze Block and the Cathaysia Block (present coordinates) (Fig. 1a; Li *et al.* 2007). The oldest rocks exposed in the Yangtze Block are the Kongling complex (marked ‘1’ in Fig. 1a) containing magmatic and metamorphic rocks as old as 3.3 Ga (Zhang *et al.* 2006a; Gao *et al.* 2011). Zheng *et al.* (2006) suggested a wide distribution of Archaean rocks in the unexposed basement of the Yangtze Block based on U–Pb and Hf isotopic data from zircon xenocrysts in Palaeozoic lamproites. Zhao *et al.* (2010) and Wang

et al. (2012) reached the same conclusion based on detrital zircon analyses of Palaeo- to Neoproterozoic sedimentary rocks in the southwestern Yangtze Block. These recent studies have proposed that the Archaean to Palaeoproterozoic basement is probably more widespread in the Yangtze Block than previously thought (Zheng *et al.* 2006; Zhang *et al.* 2006a,b; Wang *et al.* 2010, 2012).

The Kangdian area (Fig. 1b) is located near the southwestern margin of the Yangtze Block (Fig. 1a). The oldest supracrustal rocks in this area are the late Palaeo- to Mesoproterozoic meta-volcanic and meta-sedimentary rocks, termed the Dahongshan Group (Greentree & Li, 2008), the Lower Kunyang Group (Dongchuan Group; Zhao *et al.* 2010), the Hekou Group (Wu *et al.* 1990; D. F. He, unpub. Ph.D. thesis, Graduate School of the Chinese Academy of Sciences, 2009; Zhao & Zhou, 2011), the Huili Group (Yin, Sun & Zhang, 2011) and the Kunyang Group (Yin, Sun & Zhang, 2011), which occur along the Luzhijiang fault and a series of related NNE-trending faults (Fig. 1b). They consist of meta-sedimentary rocks interbedded with felsic and mafic meta-volcanic rocks (Li *et al.* 2002, 2006; Greentree *et al.* 2006; Greentree & Li, 2008; Zhao *et al.* 2010). These late Palaeo- to Mesoproterozoic rocks are overlain by a thick sequence (maximum > 9 km) of Neoproterozoic (850–540 Ma) to Permian strata consisting of clastic, carbonate and volcanic rocks (Cong, 1988; SB GMR, 1991).

The ages of the Precambrian units in the Kangdian region had been poorly constrained (see review by Wu *et al.* 1990) until the availability of precise U–Pb zircon ages in recent years. The Dahongshan Fe–Cu ore deposit is hosted in the Dahongshan Group (Zhao & Zhou, 2011). Meta-volcanic units in the Dahongshan Group have been dated at 1675 ± 8 Ma and 1681 ± 13 Ma by SHRIMP U–Pb zircon (Greentree & Li, 2008) and LA-ICP-MS U–Pb zircon (Zhao & Zhou, 2011) methods, respectively.

The Kunyang Group in the southern part of the study region (Fig. 1b) was divided into the Upper and Lower Kunyang groups (e.g. Zhao *et al.* 2010; Zhao & Zhou, 2011; Yin, Sun & Zhang, 2011). From the base upwards, the Lower Kunyang Group (Zhao *et al.* 2010; Zhao & Zhou, 2011; Yin, Sun & Zhang, 2011) includes the Yinmin, Luoxue, E'touchang and Luzhijiang formations, whereas the Upper Kunyang Group consists of the Dayingpan, Heishantou, Dalongkou and Meidang formations. Several 'Dongchuan-type' Fe–Cu ore deposits are hosted in the Lower Kunyang Group (Zhao *et al.* 2010, 2013; Zhao & Zhou, 2011). Detrital zircons from the Lower Kunyang Group yielded the youngest age of c. 1.78 Ga, and a tuff sample from the same group has a U–Pb zircon age of 1742 ± 13 Ma (Zhao *et al.* 2010). Sun *et al.* (2009) recently reported a SHRIMP zircon age of 1503 ± 17 Ma for a tuff sample from the E'touchang Formation in the Lower Kunyang Group. Thus, the Lower Kunyang Group likely formed between c. 1.7 Ga and c. 1.5 Ga. Zhao *et al.* (2010) hence interpreted the Lower Kunyang (Dongchuan)

and Dahongshan groups as stratigraphically correlatable units (Fig. 1b). In addition, a carbonaceous shale sample of the Dayingpan Formation from the Upper Kunyang Group yielded a whole-rock Pb–Pb isochron age of 1258 ± 70 Ma (Chang *et al.* 1997). A tuff layer in the Heishantou Formation of the Upper Kunyang Group has been dated at 995 ± 15 Ma and 1032 ± 9 Ma by SHRIMP U–Pb zircon dating (Greentree *et al.* 2006; Zhang *et al.* 2007). Meanwhile, detrital zircons from the Upper Kunyang Group also gave U–Pb ages as young as 960 Ma (Greentree *et al.* 2006).

The Huili Group in the northern part of the study region (Fig. 1b) is an over 10 km thick sequence of meta-clastic and meta-carbonate rocks interbedded with volcanic rocks (Wu *et al.* 1990). The group consists of the Yinmin, Luoxue, Heishan, Qinglongshan, Limahe, Fengshan and Tianbaoshan formations, from the base upwards (Wu *et al.* 1990). The meta-volcanic rocks in the Tianbaoshan Formation have TIMS U–Pb zircon and SHRIMP U–Pb zircon ages of 958 ± 16 Ma (Mou, Lin & Yu, 2003) and 1028 ± 9 Ma (Geng *et al.* 2007), respectively.

The Hekou Group in the northwestern part of the study region (Fig. 1b) is a sequence of meta-clastic and meta-carbonate rocks. This group is in fault contact with the Huili Group to the W and NE and is unconformably overlain by Sinian and Phanerozoic strata to the NW and E–SE (Fig. 2a). The Hekou Group comprises the Dayingshan, Luodang and Changchong formations from the base upwards (Fig. 2b), and has a total thickness of > 1800 m (Fig. 2b; J. Y. Zhou, unpub. Ph.D. thesis, Chengdu Univ. Technology, 2005). The Lala Fe–Cu ore deposit is hosted in the Hekou Group (D. F. He, unpub. Ph.D. thesis, Graduate School of the Chinese Academy of Sciences, 2009; Chen & Zhou, 2012). The latter is generally considered to be the equivalent of the Dahongshan Group (Wu *et al.* 1990; Zhao *et al.* 2010), as confirmed by a recent SHRIMP U–Pb zircon age of 1695 ± 20 Ma for the meta-volcanic rocks (D. F. He, unpub. Ph.D. thesis, Graduate School of the Chinese Academy of Sciences, 2009) (Figs 1, 2).

A number of relatively well-preserved mafic and felsic intrusions are exposed in the vicinity of the Hekou Town and Lala Fe–Cu ore deposit in the Hekou area. The former intruded into the Dayingshan Formation and the latter into Luodang Formation of the Hekou Group, respectively (Fig. 2a, b; SB G, 1967; D. F. He, unpub. Ph.D. thesis, Graduate School of the Chinese Academy of Sciences, 2009; Zhou *et al.* 2009; Guan *et al.* 2011; Chen & Zhou, 2012). Rocks from these mafic intrusions are slightly to strongly altered to chlorite, actinolite, albite and magnetite, but their ophitic texture and minor primary clinopyroxene are still preserved (J. Y. Zhou, unpub. Ph.D. thesis, Chengdu Univ. Technology, 2005; Chen & Zhou, 2012).

The Hekou gabbroic intrusion investigated in this paper intruded into the Dayingshan Formation of the Hekou Group near Hekou Town, SW Huili County, Sichuan province (Fig. 1b). The intrusion strikes NE–NW and dips to the SE or SW with an angle of




Figure 2. (Colour online) (a) A simplified geologic map of the Hekou area, Sichuan Province, SW China; (b) a simplified stratigraphic column of the Hekou Group. Modified from the Chinese Huili 1:200 000 geological map (SB G, 1967).

~30–45°. The intrusion is the biggest mafic intrusion in the Hekou area and is up to 4–5 km long with a thickness between 1 and 2 km, covering an area of about 6 km² (Fig. 2a). These rocks are dark-grey to black, fine to coarse grained, with massive and subhedral granular and ophitic textures. The coarse-grained gabbroic rocks are in the central zone (CZ) of the intrusion, and are moderately to strongly altered to chlorite, actinolite and albite. However, the medium- to fine-grained gabbroic rocks are usually observed in the outer zone (OZ) of the intrusion, and are relatively fresh to slightly altered. These rocks mainly consist of plagioclase (50–55 vol. % by

volume) and clinopyroxene (40–45 vol. %), and minor subordinate orthopyroxene, Fe–Ti oxides and apatite. Clinopyroxenes in the gabbros from the OZ commonly exhibit a medium- to fine-grained texture. Electron microprobe analyses indicate clinopyroxenes from the gabbros are $\text{Wo}_{38-49}\text{En}_{34-46}\text{Fs}_{11-22}$, mostly belonging to augite (online Supplementary Material available at <http://journals.cambridge.org/geo>).

3. Sampling and analytical methods

Nineteen samples were collected from the CZ (samples HK1101 to HK1103) and the OZ (samples HK1104 to

HK1211) of the gabbroic intrusion in the Hekou area (Fig. 2a, b).

Zircons from samples HK1103 and HK1107 were separated using conventional heavy liquid and magnetic techniques, then by handpicking under binocular microscopes. They were mounted in an epoxy resin disc, and polished and coated with gold film. Zircons were documented with transmitted and reflected light micrographs as well as cathodoluminescence (CL) images to reveal their external and internal structures. U–Pb zircon dating was conducted using a Cameca IMS 1280 ion microprobe (SIMS) at the Institute of Geology and Geophysics, the Chinese Academy of Sciences (CAS) in Beijing. Details of the analytical procedures for U–Pb zircon dating can be found in Li *et al.* (2009, 2010). Oxygen flooding that introduces oxygen into the sample chamber was used during the analyses, which not only enhances Pb^{+} ion yield by a factor of 2 and 7 for zircon and baddeleyite (Li *et al.* 2009), respectively, but also depresses the U–Pb orientation effect (Wingate & Compston, 2000) down to $\sim 2\%$ (Li *et al.* 2010). The uncertainties in ages are cited as 1σ , and the weighted mean ages are quoted at the 95% confidence interval (2σ). The SIMS U–Pb zircon ages are presented in Table 1.

Chemical compositions of clinopyroxenes from the gabbroic intrusion were determined by wavelength-dispersion X-ray emission spectrometry using an EPMA-1600 electron microprobe at the State Key Laboratory of Ore Deposit Geochemistry (SKL OD G), Institute of Geochemistry, Chinese Academy of Sciences (IGCAS). Major-element compositions of whole rocks were determined using X-ray fluorescence spectrometers (XRF) at the State Key Laboratory of Isotope Geochemistry, Guangzhou Institute of Geochemistry of the CAS. The analytical precision was better than 5%. Trace elements in whole rocks were analysed using a Perkin-Elmer Sciex ELAN DRC–e ICP-MS at the SKL OD G, IGCAS. The powdered samples (50 mg) were dissolved with $HF + HNO_3$ mixture in high-pressure Teflon bombs at $\sim 190^\circ C$ for 48 hours (Qi, Hu & Gregoire, 2000). Rh was used as an internal standard to monitor signal drift during measurement. The international standards GBP G-1, OU-6 and the Chinese National standards GSR-1 and GSR-3, were used for analytical quality control. The analytical precision is generally better than 10% for trace elements.

Samples for Nd isotopic analyses were dissolved in an acidic mixture of HF , HNO_3 and $HClO_4$ in Teflon bombs. Sm and Nd were separated by conventional cation-exchange techniques at the SKL OD G. The Nd isotopic measurements were performed on a multicollector ICP-MS (MC-ICP-MS) at the Key Laboratory of Orogenic Belts and Crustal Evolution, School of Earth and Space Sciences, Peking University. The mass fractionation corrections for Nd isotopic ratios are based on $^{146}Nd/^{144}Nd = 0.7219$. The $^{143}Nd/^{144}Nd$ ratios of the JNDI-1 Nd standard solution and the US GS standard rock BCR-2 were determined to be 0.512119 ± 14 (2σ) and 0.512629 ± 16 (2σ), respectively.




Figure 3. (Colour online) Representative CL images of zircons from (a) sample HK1003 from the CZ; and (b) sample HK1007 from the OZ of the gabbroic rocks from the Hekou intrusion.

4. Results

4.a. U–Pb zircon geochronology

4.a.1. Sample HK1103 ($26^\circ 16' 12.7'' N$, $102^\circ 01' 04.6'' E$)

Zircons extracted from sample HK1103 in the CZ are mostly clear and up to 50–100 μm in length with length to width ratios between 1:1 and 1.5:1. Most zircons are simple prismatic crystals without obvious zoning under CL (Fig. 3a). Fourteen analyses were conducted on 14 zircons. Uranium and thorium concentrations are variable, with U = 213–1263 ppm, Th = 29–263 ppm.

Table 1. Cameca SIMS U–Pb zircon isotopic analyses for the gabbroic rocks from the Hekou intrusion, SW China

Spot	U (ppm)	Th (ppm)	Th/U	²⁰⁶ Pb/ ²⁰⁴ Pb measured	<i>f</i> ₂₀₆ (%)	Isotopic ratio						Age/Ma	
						²⁰⁷ Pb / ²⁰⁶ Pb	± 1σ (%)	²⁰⁷ Pb / ²³⁵ U	± 1σ (%)	²⁰⁶ Pb / ²³⁸ U	± 1σ (%)	²⁰⁶ Pb / ²³⁸ U ± 1σ	²⁰⁷ Pb / ²⁰⁶ Pb ± 1σ
HK1103 (26° 16′ 12.7 N, 102° 01′ 04.6 E)													
1	213	462	2.169	55680	0.03	0.10679	0.70	4.51721	1.67	0.3068	1.52	1725 ± 23	1745 ± 13
2	423	419	0.991	29413	0.06	0.10685	0.40	4.28050	1.56	0.2906	1.50	1644 ± 22	1746 ± 7
3	1263	820	0.649	34369	0.05	0.10645	0.29	4.54819	1.53	0.3099	1.50	1740 ± 23	1740 ± 5
4	761	2879	3.786	135897	0.01	0.10682	0.31	4.47788	1.54	0.3040	1.51	1711 ± 23	1746 ± 6




Figure 4. SIMS U–Pb Concordia diagrams of (a) sample HK1003 from the CZ; and (b) sample HK1007 from the OZ of the gabbroic rocks from the Hekou intrusion.

on CL images, and are 50–200 μm long with a length to width ratio of between 1:1 and 2:1 (Fig. 3b). Seventeen Cameca SIMS analyses were completed on 17 zircons with U concentration between 302 and 1533 ppm, Th between 388 and 2527 ppm, and Th/U ratios between 0.96 (




Figure 5. Diagrams of Y, Nb, Th, La, V, Sr, Pb, Rb and Ba v. Zr to evaluate the mobility of these elements in the gabbroic rocks from the Hekou intrusion during alteration. Solid round symbols – the central zone (CZ) of the gabbroic intrusion; Open round symbols – the outer zone (OZ) of the gabbroic intrusion.

altered to varying degrees judging from thin-section examinations. Nineteen samples from the Hekou gabbroic intrusion have been selected for major- and trace-element analyses (Table 2). The loss of ignition (LOI) values for the gabbroic rocks range from 0.55 to 2.69%. Thus, the effects of alteration on the chemical compositions of these rocks need to be evaluated. Zirconium in mafic intrusive rocks is generally considered to be the most immobile during low- to medium-grade alteration (e.g. Wood, Joron & Treuil, 1979; Gibson *et al.* 1982). Therefore, bivariate plots of Zr against selected trace elements have been used for evaluating the mobilities of such elements during alteration (e.g. Polat, Hofmann & Roseng, 2002). A number of elements with different geochemical behaviours, including Y, Nb, Th, La, V, Sr, Rb, Rb and Ba in the gabbroic rocks, are plotted against Zr to evaluate their mobility during alteration. Except for Pb, Rb and Ba, the other elements overall correlated tightly with Zr, indicating that these elements are essentially immobile during alteration (Fig. 5). Therefore, only the immobile elements are used for geochemical classification and petrogenetic discussions in this study. Meanwhile, the sums of major-element oxides for all samples in this study are recalculated to 100% volatile free.

4.b.2. Major- and trace-element geochemistry

The rocks in the CZ of the Hekou gabbroic intrusion show fairly homogeneous major- and trace-element compositions, with Mg no. from 61.5 to 62.3, SiO₂ from 48.2 to 49.0%, TiO₂ from 0.81 to 0.92%, MgO ~8.9%, Fe₂O₃ from 10.7 to 11.0%, CaO from 10.1 to 11.2%, Al₂O₃ from 15.9 to 16.7%, Zr from 34.7

to 38.5 ppm, Cr from 312 to 321 ppm, and Ni from 141 to 171 ppm (Table 2; Fig. 6). In contrast, the rocks from the OZ display slightly evolved compositions, with relatively lower Mg no. (54.3–59.3), MgO (6.9–8.4%), Al₂O₃ (14.1–16.4%), Cr (203–356 ppm) and Ni (74.6–113 ppm) contents, and higher SiO₂ (48.7–50.3%), CaO (10.4–13.2%) and Zr (40.9–56.5 ppm) concentrations than those of the rocks from the CZ of the intrusion. Meanwhile, TiO₂ (0.75–0.92%) and Fe₂O₃ (48.7–50.3%) contents of these rocks are similar to those of the rocks from the CZ (Table 2; Fig. 6). In the Fenner diagrams (Fig. 6), with decreasing MgO contents, Cr and Ni contents of the rocks from the OZ obviously decrease, and Fe₂O₃ contents slightly decrease, whereas SiO₂, TiO₂, CaO and Al₂O₃ contents slightly increase, and Zr contents obviously increase with decreasing MgO contents (Fig. 6). The Ti/Y ratios of the gabbroic rocks are 467–550 and 325–399 for the CZ and OZ, respectively. On the TAS rock classification diagram, all the gabbroic rocks from the intrusion are plotted in the subalkaline field (Fig. 7). Therefore, the gabbroic rocks from the Hekou intrusion were related to subalkaline basaltic affinities.

The gabbroic rocks from the Hekou intrusion have low total REE contents (REE = 29.3–40.2 ppm) (Fig. 8a). These rocks in the CZ of the intrusion show slightly LREE-enriched and HREE-depleted patterns ($(La_N = 14–18, (La/Yb)_N = 3.1–4.0, (La/Sm)_N = 1.8–1.9, (Gd/Yb)_N = 1.4–1.7$; subscript N denotes chondrite normalized) (Boynton, 1984), and slightly positive Eu anomalies ($Eu/Eu = 1.2–1.3$) (Fig. 8a). Compared with the rocks from the CZ of the intrusion, the OZ rocks have relatively flatter LREE and HREE patterns, with $La_N = 14–18, (La/Yb)_N = 2.1–2.7,$

Table 2. Major-element (in wt %) and trace-element (in ppm) data for the gabbroic rocks from the Hekou intrusion

Rock type Sample no.	Gabbro HK1101	Gabbro HK1102	Gabbro HK1103	Gabbro HK1104	Gabbro HK1105	Gabbro HK1106	Gabbro HK1107	Gabbro HK1108
SiO ₂	47.55	47.41	48.03	49.10	49.14	48.08	49.27	48.57
TiO ₂	0.79	0.83	0.90	0.84	0.86	0.81	0.86	0.88
Al ₂ O ₃	15.64	15.84	16.39	14.11	14.25	14.42	14.79	15.34
Fe ₂ O ₃	10.46	10.81	10.55	11.21	10.68	11.31	11.06	10.57
MnO	0.09	0.10	0.12	0.22	0.20	0.20	0.19	0.21
MgO	8.73	8.70	8.71	7.50	7.64	7.77	7.38	7.00

Table 2. Continued.

Rock type Sample no.	Gabbro HK1201	Gabbro HK1202	Gabbro HK1203	Gabbro HK1204	Gabbro HK1205	Gabbro HK1206	Gabbro HK1207	Gabbro HK1208	Gabbro HK1209	Gabbro HK1210	Gabbro HK1211
SiO ₂	48.78	48.88	48.45	48.26	48.81	48.94	48.75	48.71	48.35	48.68	47.57
TiO ₂	0.83	0.86	0.73	0.74	0.81	0.90	0.80	0.84	0.79	0.87	0.87
Al ₂ O ₃	15.39	15.83	15.80	15.77	16.04	14.64	14.69	15.54	14.93	13.75	14.42
Fe ₂ O ₃	10.51	10.34	9.95	10.38	9.85	11.44	10.76	10.25	10.83	11.43	11.19
MnO	0.21	0.18	0.21	0.20	0.21	0.21	0.22	0.20	0.26	0.19	0.21
MgO	7.54	6.84	7.20	7.17	6.80	6.85	7.91	7.31	7.81	8.19	8.18
CaO	10.09	13.05	12.02	11.69	12.33	12.76	12.39	12.01	11.58	11.92	12.90
Na ₂ O	2.77	2.20	2.69	2.83	2.37	1.58	1.84	2.30	2.42	1.90	2.05
K ₂ O	0.87	0.31	0.19	0.17	0.39	0.19	0.30	0.54	0.78	0.87	0.23
P ₂ O ₅	0.05	0.07	0.06	0.06	0.07	0.07	0.06	0.07	0.06	0.06	0.07
LOI	2.69	1.19	2.32	2.09	2.12	2.18	1.85	1.59	1.79	1.52	2.12
Total	99.73	99.75	99.62	99.36	99.80	99.76	99.57	99.36	99.60	99.38	99.81
Mg no.	58.7	56.7	58.9	57.8	57.8	54.3	59.3	58.5	58.8	58.7	59.1
Sc	38.7	36.9	35.3	36.1	33.7	37.0	37.9	37.0	37.5	40.7	39.2
V	209	187	183	224	224	180	244	241	240	201	256
Cr	265	228	231	218	203	255	270	270	278	356	355
Co	48.7	46.3	38.6	40.1	38.8	42.1	47.0	47.1	47.9	52.0	42.8
Ni	99.2	88.2	78.3	75.6	74.6	80.1	91.0	89.8	92.5	101.4	91.9
Cu	108	93.3	108	116	83.0	83.7	110	102	98.9	94.2	72.1
Zn	97.0	106	248	374	127	86.5	135	150	214	104	94.6
Ga	13.8	15.8	15.0	15.0	15.5	16.9	13.7	15.3	14.7	14.3	14.6
Rb	59.7	12.8	6.96	6.27	23.5	2.94	16.1	31.4	48.8	43.8	4.92
Sr	219	181	266	244	290	245	203	199	246	177	183
Y	13.0	14.3	13.3	14.0	14.5	14.6	13.5	14.3	13.7	14.6	15.5
Zr	40.9	50.6	44.6	49.8	53.1	56.5	44.1	48.7	47.0	46.2	47.8
Nb	3.17	3.95	3.35	3.54	3.99	4.26	3.26	3.79	3.42	3.53	3.65
Cs	0.396	0.336	0.050	0.046	0.184	0.088	0.141	0.387	0.705	0.324	0.025
Ba	277	131	106	108	329	109	214	163	494	264	121
La	4.34	5.20	4.83	5.04	5.63	5.67	4.45	5.18	4.44	4.73	5.24
Ce	10.0	11.9	11.1	11.5	12.7	12.6	10.3	11.8	10.4	10.7	11.6
Pr	1.40	1.68	1.56	1.61	1.75	1.71	1.45	1.65	1.50	1.56	1.68
Nd	6.19	7.40	6.74	7.14	7.74	7.67	6.44	7.32	6.80	6.94	7.75
Sm	1.74	2.13	1.87	2.04	2.12	2.13	1.86	2.09	1.94	2.06	2.13
Eu	0.774	0.833	0.804	0.843	0.910	0.930	0.739	0.798	0.906	0.846	0.808
Gd	2.08	2.21	2.17	2.23	2.28	2.33	2.04	2.17	2.20	2.23	2.39
Tb	0.384	0.431	0.402	0.428	0.436	0.435	0.401	0.419	0.405	0.436	0.443
Dy	2.33	2.59	2.38	2.49	2.54	2.61	2.45	2.53	2.49	2.65	2.75
Ho	0.529	0.587	0.550	0.584	0.576	0.598	0.559	0.596	0.569	0.593	0.624
Er	1.40	1.52	1.44	1.48	1.52	1.58	1.44	1.54	1.51	1.58	1.60
Tm	0.206	0.224	0.200	0.210	0.217	0.226	0.219	0.201	0.215	0.226	0.233
Yb	1.28	1.42	1.34	1.42	1.41	1.49	1.28	1.42	1.39	1.49	1.50
Lu	0.192	0.222	0.188	0.202	0.211	0.217	0.202	0.199	0.201	0.218	0.222
Hf	1.27	1.51	1.39	1.58	1.56	1.72	1.38	1.51	1.45	1.44	1.49
Ta	0.195	0.255	0.215	0.230	0.339	0.269	0.217	0.237	0.217	0.223	0.233
Pb	8.00	10.5	24.2	10.8	12.7	3.49	8.13	8.59	54.7	6.45	4.93
Th	0.635	0.753	0.698	0.743	0.794	0.886	0.669	0.746	0.684	0.676	0.716
U	0.147	0.168	0.156	0.177	0.185	0.180	0.141	0.175	0.160	0.153	0.171

Mg no. = 100 molar MgO/(Mg + FeO), assuming FeO_T = 0.9 × Fe₂O₃. Total iron as FeO_T. LOI = loss on ignition.




Table 3. Sm–Nd isotopic compositions for the gabbroic rocks from the Hekou intrusion

Sample	Sm (ppm)	Nd (ppm)	$^{147}\text{Sm}/^{144}\text{Nd}$	$^{143}\text{Nd}/^{144}\text{Nd}$ (2 σ)	$(^{143}\text{Nd}/^{144}\text{Nd})_i$	$\varepsilon_{\text{Nd}}(\text{T})$
HK1101	1.916	7.430	0.1559	0.512208 (9)	0.510428	0.70
HK1102	1.517	5.769	0.1590	0.512327 (17)	0.510511	2.33
HK1103	1.794	6.850	0.1583	0.512273 (16)	0.510466	1.45
HK1106	1.917	6.538	0.1772	0.512400 (20)	0.510377	−0.30
HK1206	2.128	7.667	0.1678	0.512321 (20)	0.510404	0.24
HK1207	1.856	6.443	0.1741	0.512386 (18)	0.510398	0.12

Chondrite uniform reservoir (CHUR) values ($^{147}\text{Sm}/^{144}\text{Nd} = 0.1967$, $^{143}\text{Nd}/^{144}\text{Nd} = 0.512638$) are used for the calculation. $\lambda_{\text{Sm}} = 6.54 \times 10^{-12} \text{ year}^{-1}$ (Lugmair & Harti, 1978). The $(^{143}\text{Nd}/^{144}\text{Nd})_i$ and $\varepsilon_{\text{Nd}}(\text{T})$ of these rocks were calculated using the age of 1736 Ma.




Figure 7. Rock classification plot of SiO_2 v. $(\text{Na}_2\text{O} + \text{K}_2\text{O})$ (Cox, Bell & Pankhurst, 1979) for the gabbroic rocks from the Hekou intrusion. Symbols as in Figure 5.

5. Discussion

5.a. The petrogenesis of the gabbroic intrusion

5.a.1. Fractional crystallization

The rocks from the CZ of the Hekou gabbroic intrusion show fairly homogeneous major oxides and Mg no. values. In contrast, the gabbroic rocks from the OZ of the intrusion exhibit slightly variable MgO concentrations and Mg no. values, suggesting that they had undergone small degrees of fractional crystallization (Fig. 6). A slight increase of TiO_2 with decreasing MgO contents for these rocks from the OZ indicates that TiO_2 was incompatible in the crystallizing phases and therefore titanomagnetite did not appear on the liquidus during crystallization of plagioclase and clinopyroxene in these systems. The slightly positive correlations of Fe_2O_3 versus MgO , and the obviously positive correlations of Cr and Ni versus MgO imply that they may undergo some degrees of fractionation of orthopyroxene.

The negative correlation of Al_2O_3 v. MgO in the rocks from the OZ suggests that they underwent insignificant plagioclase fractionation (Fig. 6; Pik *et al.* 1998). The gabbroic rocks from the Hekou intrusion have relatively high CaO contents (10.1–13.2%) and low REE contents (29.3–40.2 ppm), which are consistent with the features of the mafic cumulate rocks.

Moreover, all the rocks from the intrusion display a slightly negative correlation in the CaO v. MgO diagram (Fig. 6), implying clinopyroxene is a major fractionated phase for these rocks.

In summary, clinopyroxene and minor amounts of orthopyroxene are the important fractionation phases for the rocks from the OZ of the gabbroic intrusion.

5.a.2. Crustal contamination

The gabbroic rocks from the CZ of the intrusion display significant Th depletion and insignificant Nb and Ta depletion relative to La. In contrast, the gabbroic rocks in the OZ are slightly enriched in Th and depleted in Nb and Ta (Fig. 8b). As shown in the $\varepsilon_{\text{Nd}}(\text{T})$ –Nb/La plot (Fig. 9a; Paces & Bell, 1989) and the Nb/Th–Nb/La plot (Fig. 9b), the less-contaminated rocks from the CZ have relatively higher $\varepsilon_{\text{Nd}}(\text{T})$ values and Nb–Ta contents, and lower Th contents than those of the more contaminated rocks from the OZ. In view of these geochemical and isotopic correlations, the crustal contaminants must mainly have caused low $\varepsilon_{\text{Nd}}(\text{T})$ values, and Nb/La and Nb/Th ratios, indicating sialic crust components in the OZ rocks. Furthermore, the continental crust is typically depleted in Nb and Ta (Rudnick & Fountain, 1995), and the upper continental crust is enriched in La and Th while the lower continental crust is not usually enriched in Th (Barth, McDonough & Rudnick, 2000). As a result, the crustal contaminant, which is enriched in Th and La, and depleted in Nb and Ta, was potentially derived from the upper crust rather than from the lower crust (Fig. 9c; Ingle *et al.* 2002). It is therefore suggested that the magmas parental to the gabbroic rocks assimilated the upper continental crust.

5.a.3. Nature of the parental magma and mantle source characteristics

Although the magmas parental to the Hekou gabbroic intrusion were subjected to variable degrees of crustal contamination, the rocks from the CZ exhibit obvious Th depletion and insignificant Nb, Ta and Ti depletions relative to La (Fig. 8b). However, the trace elemental features of the gabbroic rocks are different from those of typical E-MORB because: (1) all the gabbroic rocks are cumulate rocks and exhibit more obvious LREE-enriched and HREE-depleted patterns (Fig. 8a, b); (2)




Figure 8. (Colour online) (a) Chondrite-normalized REE patterns; and (b) primitive-mantle normalized incompatible trace-element multi-element plot for the gabbroic rocks from the Hekou intrusion. The normalization values of chondrite are from Boynton (1984). The normalization values of primitive mantle are from Sun & McDonough (1989). Symbols as in Figure 5. OIB – ocean island basalt; E-MORB – enriched mid-ocean ridge basalt; N-MORB – normal mid-ocean ridge basalt.

the magmas parental to the gabbroic rocks had been assimilated by sialic crust. This makes it unlikely that the gabbroic intrusion was formed in an oceanic setting where sialic crust is normally absent. In contrast, the trace elemental features of the less-contaminated gabbroic samples are very similar to the intraplate basaltic rocks in CFB and OIB provinces (Sun & McDonough, 1989). For example, the less-contaminated rocks of the CZ have high Ti/Y (467–550) and low La/Nb (0.93–1.32) and La/Ta ratios (14.3–19.3), which are also similar to those of OIB (Ti/Y > 410, La/Nb < 1.5 and La/Ta < 22) (Table 2; Ormerod *et al.* 1988; Sun & McDonough, 1989; Lightfoot *et al.* 1993; Garland, Turne & Hawkesworth, 1996; Ewart *et al.* 1998).

Based on the aforementioned results, the magmas parental to the gabbroic rocks display a subalkaline basaltic affinity. In combination with the Nd isotopic features ($\epsilon_{\text{Nd}}(\text{T}) = +0.70$ to $+2.3$) of the less-contaminated rocks from the CZ, the parental magmas for the Hekou gabbroic rocks were proposed to be generated dominantly by melting of a slightly depleted mantle source. The lack of Nb–Ta depletion and the high-Ti nature of the magma parental to the less-contaminated rocks suggest the sub-continental lithospheric mantle (SCLM) was not the geological reservoir that contributed the trace-element and Nd isotope inventory to the Hekou gabbroic intrusion. In general, low ratios of LREE/HREE in mafic rocks reflect a melting regime dominated by relatively large melting fractions and/or having spinel as the predominant residual phase, whereas high LREE/HREE ratios are indicative of smaller melting fractions and/or garnet control

(Deniel, 1998; Xu *et al.* 2001). Therefore, the samples from the Hekou gabbroic intrusion, exhibiting a slightly LREE-enriched and HREE-depleted pattern, were a




Figure 9. Plots of (a) $\epsilon_{\text{Nd}}(T)$ v. Nb/La; (b) Nb/Th v. Nb/La; and (c) $(\text{Th}/\text{Ta})_{\text{PM}}$ v. $(\text{La}/\text{Ta})_{\text{PM}}$ (Ingle *et al.* 2002) for the gabbroic rocks from the Hekou intrusion. Symbols as in Figure 5. UCC – upper continental crust; LCC – lower continental crust.

were generally intensely altered and did not preserve their original chemical characteristics. Therefore, a major problem with better understanding these deposits is the lack of precise elemental and isotopic data for these meta-volcanic rocks and mafic intrusions, which hampers the understanding of the geodynamic setting of the ore mineralization.

The Hekou gabbroic intrusion has a SIMS U–Pb zircon age of *c.* 1736 Ma reported in this study, suggesting that the depositional ages of the Hekou Group were slightly older than the ages of the intrusion (Chen, Zhou & Zhao, 2013). Moreover, the crystallization ages of the gabbroic intrusion are similar to that of the dolerite dykes that intruded into the Dahongshan Group in the Dahongshan Fe–Cu ore deposit (Zhao & Zhou, 2011)




Figure 10. Discrimination diagram of Al_2 (percentage of tetrahedral sites occupied by Al) v. TiO_2 in clinopyroxenes (Loucks, 1990) for the gabbroic rocks from the Hekou intrusion. Symbols as in Figure 5.

and into the Lower Kunyang Group in the Yinmin Fe–Cu ore deposit (Zhao *et al.* 2010, 2013), indicating that significant mafic magmatism occurred at *c.* 1.7 Ga in the southwestern Yangtze Block. The meta-volcanic rocks from the Dahongshan Group and the Yinmin Formation in the Lower Kunyang Group were also formed slightly before or almost synchronously with the *c.* 1.7 Ga mafic dykes and intrusions (Greentree & Li, 2008; D. F. He, unpub. Ph.D. thesis, Graduate School of the Chinese Academy of Sciences, 2009; Zhao & Zhou, 2011; Zhao *et al.* 2010, 2013). Therefore, the late Palaeoproterozoic to Mesoproterozoic volcanic and sedimentary basins, termed the Dahongshan, Hekou and Lower Kunyang groups, were likely formed almost synchronously and slightly prior to *c.* 1.7 Ga.

As discussed above, the least-contaminated gabbroic rocks from the CZ exhibit characteristic patterns of incompatible trace elements similar to those of OIB and CFB (Fig. 8b). As shown in Figure 10, the clinopyroxene crystals from the Hekou intrusion show the characteristics of gabbroic cumulate rocks formed in a continental rift system. Magmatism associated with lithosphere rifting coupled with asthenosphere upwelling can explain why the intrusion has isotopic compositions similar to those of magmas derived from the asthenospheric mantle. Therefore, the gabbroic intrusion intruded into the Hekou Group is best interpreted as a product of a continental rift environment.

In the Kangdian region, the Yinmin Formation of the Lower Kunyang Group with a U–Pb zircon age of 1742 ± 13 Ma has been considered to have formed in a continental rift basin (Hua, 1990; Wu *et al.* 1990; Zhao *et al.* 2010). Meanwhile, alkali basaltic tuff is present in the Yinmin Formation (Wu *et al.* 1990; Greentree *et al.* 2006). Moreover, the $\epsilon_{\text{Hf}}(t)$ values of igneous zircons from the tuff and ~ 1.70 to 1.65 Ga dolerite dykes that intruded into the Yinmin Formation suggest they were derived from a depleted mantle source. These dolerite dykes show no anomalies of Nb and Ta on the

primitive-mantle normalized multi-element plot, and fall into the field of within-plate basalt (Zhao *et al.* 2010, 2013; Zhao & Zhou, 2011). These sedimentary records combined with the elemental and isotopic data from these igneous rocks support the interpretation that the Yinmin Formation of the Lower Kunyang Group was formed in a continental rift setting at ~1.7 Ga. It is therefore suggested that the c. 1.7 Ga Dahongshan, Hekou and Lower Kunyang groups were likely formed in similar continental rift basins in the southwestern Yangtze Block.

Collisional events at 2.1 to 1.8 Ga have been recorded in a number of Precambrian cratons and are thought to be related to the assembly of the Palaeoproterozoic supercontinent Columbia (Rogers & Santosh, 2002; Zhao *et al.* 2002a). More recently, it was suggested that even the 1.84–1.75 Ga mafic dyke swarms and 1.75–1.68 Ga anorthosite–mangerite–alkali granite and rapakivi granite suites (AMGRS) in the North China Craton (NCC) may have resulted from the fragmentation of the Columbia supercontinent (e.g. Zhao *et al.* 2002b; Peng, Zhai & Guo, 2006; Peng *et al.* 2007, 2008; Hou *et al.* 2008a,b). The late Palaeo- to Mesoproterozoic anorogenic magmatism associated with the break-up of the Columbia supercontinent generally took place during intraplate continental rifting, triggered mostly by mantle plumes (Rogers & Santosh, 2002; Zhao *et al.* 2002a, 2004; Peng *et al.* 2008; Hou *et al.* 2008a,b; Evans & Mitchell, 2011).

In the traditional reconstruction models (Rogers & Santosh, 2002; Zhao *et al.* 2002a), the Yangtze Block was not included in the Columbia supercontinent owing to the lack of geological and geochronological data for that time. However, recent studies reveal that the Yangtze Block may have been involved in this global Palaeoproterozoic orogenic event (Zhang *et al.* 2006b; Sun *et al.* 2008; Wu *et al.* 2008; Peng *et al.* 2009, 2012) and late Palaeo- to Mesoproterozoic anorogenic magmatism (Zhao *et al.* 2010, 2013; Chen, Zhou & Zhao, 2013). In this study, the presence of the c. 1.7 Ga Hekou gabbroic intrusion in the southwestern Yangtze Block is contemporaneous with the break-up of the Columbia supercontinent, which provides further evidence to support the idea that the Yangtze Block was likely a part of the supercontinent.

It is difficult to tell the precise position of the Yangtze Block within the Columbia supercontinent without a large number of reliable palaeomagnetic poles. However, on the basis of similar age spectra of detrital zircons, the geochemical characteristics of the Palaeoproterozoic strata, comparisons of rift-related sediments and ~1.7 Ga within-plate mafic magmatism (such as the Hekou intrusion reported in this paper) among different blocks in the Columbia supercontinent, the Yangtze Block is considered to possibly link with the North Australian Craton during Palaeoproterozoic time (Wang *et al.* 2012; Chen, Zhou & Zhao, 2013; Chen *et al.* 2013; Chen, Zhang & Song, 2013; Wang & Zhou, 2014), as reconstructed by Zhao *et al.* (2002a, 2004). Moreover, the Yangtze Block, the NW India and

the North Australian Craton were perhaps neighbours located in a spatial association within the Columbia supercontinent, based on the similar mineralization ages and styles of the Palaeoproterozoic Fe–Cu ore deposits on these cratons (Zhou *et al.* 2014).

6. Conclusions

(1) The age of crystallization for the Hekou gabbroic intrusion is ~1.7 Ga, as demonstrated by the upper intercept ages of 1735 ± 6.5 Ma and 1736 ± 4.0 Ma for the rocks from the CZ and the OZ, respectively.

(2) The Hekou gabbroic intrusion experienced slight fractional crystallization from a parental magma, which was potentially generated by melting of a slightly depleted asthenospheric mantle source in a continental rift environment plus varying degrees of crustal contamination.

(3) The Yangtze Block was likely a fragment of the Palaeo- to Mesoproterozoic supercontinent Columbia, and the c. 1.7 Ga Hekou gabbroic intrusion from the southwestern Yangtze Block was likely formed in an anorogenic continental rift that contributed to the break-up of the Columbia supercontinent.

Acknowledgements. We appreciate Q. L. Li, Y. Liu, G. Q. Tang and H. Tao for their assistance with SIMS dating, Y. Liu for major-element analyses by XRF, J. Hu, G. P. Bao and Y. Huang for trace-element analyses by ICP-MS, B. L. Wang for Nd isotope analyses by MC-ICP-MS, and G. F. Zhou, and W. Q. Zheng, for electron microprobe analysis. This paper has benefited from constructive comments of the editor and anonymous reviewers. This work was funded by the NSFC (Grants 41273049, 41572074 and 41073043) and the State Key Laboratory of Ore Deposit Geochemistry (12th Five-Year Plan: SKLO G-ZY125-06).

Supplementary material

To view supplementary material for this article, please visit <http://dx.doi.org/10.1017/S0016756815001119>.

References

- BARTH, M. G., McDONOUGH, W. F. & RUDNICK, R. L. 2000. Tracking the budget of Nb and Ta in the continental crust. *Chemical Geology* **165**, 197–213.
- BOYNTON, W. V. 1984. Geochemistry of the rare earth elements: meteorite studies. In *Rare Earth Element Geochemistry* (ed. P. Henderson), pp. 63–114. Amsterdam: Elsevier.
- CHANG, X., ZHU, B., SUN, D., QIU, H. & ZOU, R. 1997. Isotope geochemistry study of Dongchuan copper deposits in Middle Yunnan Province, SW China: stratigraphic chronology and application of geochemical exploration by lead isotopes. *Geochimica* **26**, 32–8 (in Chinese with English abstract).
- CHEN, Z. C., LIN, W., FAURE, M., LEPVRIER, C., CHU, Y. & WANG, Q. C. 2013. Geochronological constraint of early Mesozoic tectonic event at Northeast Vietnam. *Acta Petrologica Sinica* **29**, 1825–40.
- CHEN, L., ZHANG, Z. & SONG, H. 2013. Weak depth and along-strike variations in stretching from a

- multi-episodic finite stretching model: evidence for uniform pure-shear extension in the opening of the South China Sea. *Journal of Asian Earth Sciences* **78**, 358–70.
- CHEN, W. T. & ZHOU, M. F. 2012. Paragenesis, stable isotopes, and molybdenite Re–Os isotope age of the Lala iron–copper deposit, Southwest China. *Economic Geology* **107**, 459–80.
- CHEN, W. T., ZHOU, M. F. & ZHAO, X. F. 2013. Late Paleoproterozoic sedimentary and mafic rocks in the Hekou area, SW China: implication for the reconstruction of the Yangtze Block in Columbia. *Precambrian Research* **231**, 61–77.
- CONG, B. L. (ed.) 1988. *Formation and Evolution of Panxi Paleo-Rift*. Beijing: Science Press, 424 pp (in Chinese with English abstract).
- COX, K. G., BELL, J. D. & PANKHURST, R. J. 1979. *The Interpretation of Igneous Rocks*. London: Allen and Unwin, 450 pp.
- DENIEL, C. 1998. Geochemical and isotopic (Sr, Nd, Pb) evidence for plume–lithosphere interactions in the genesis of Grande Comore magmas (Indian Ocean). *Chemical Geology* **144**, 281–303.
- EVANS, D. A. D. & MITCHELL, R. N. 2011. Assembly and breakup of the core of Paleoproterozoic–Mesoproterozoic supercontinent Nuna. *Geology* **39**, 443–6.
- EWART, A., MILNER, S. C., ARMSTRONG, R. A. & DUNCAN, A. R. 1998. Etendeka volcanism of the Goboboseb Mountains and Messum igneous complex, Namibia. Part I: geochemical evidence of early Cretaceous Tristan Plume melts and the role of crustal contamination in the Parana–Etendeka CFB. *Journal of Petrology* **39**, 191–225.
- FAN, H. P., ZHU, W. G., LI, Z. X., ZHONG, H., BAI, Z. J., HE, D. F., CHEN, C. J. & CHAO, C. Y. 2013. Ca. 1.5 Ga mafic magmatism in South China during the break-up of the supercontinent Nuna/Columbia: the Zhuqing Fe–Ti–V oxide ore-bearing mafic intrusions in western Yangtze Block. *Lithos* **168–169**, 85–98.
- GAO, S., YANG, J., ZHOU, L., LI, M., HU, Z., GUO, J., YUAN, H., GONG, H., XIAO, G. & WEI, J. 2011. Age and growth of the Archean Kongling terrain, South China, with emphasis on 3.3 Ga granitoid gneisses. *American Journal of Science* **311**, 153–82.
- GARLAND, F., TURNE, S. & HAWKESWORTH, C. 1996. Shifts in the source of the Paraná basalts through time. *Lithos* **37**, 223–43.
- GENG, Y., YANG, C., DU, L., WANG, X., REN, L. & ZHOU, X. 2007. Chronology and tectonic environment of the Tianbaoshan formation: new evidence from zircon SHRIMP U–Pb age and geochemistry. *Geological Review* **53**, 556–63 (in Chinese with English abstract).
- GIBSON, S. A., KIRKPATRICK, R. J., EMMERMAN, R., SCHMINCKE, P. H., PRITCHARD, G., OKAY, P. J., HORPE, R. S. & MARRINER, G. F. 1982. The trace element composition of the lavas and dykes from a 3 km vertical section through a lava pile of Eastern Iceland. *Journal of Geophysical Research* **87**, 6532–46.
- GREENTREE, M. R. & LI, Z. X. 2008. The oldest known rocks in south-western China: SHRIMP U–Pb magmatic crystallization age and detrital provenance analysis of the Paleoproterozoic Dahongshan Group. *Journal of Asian Earth Sciences* **33**, 289–302.
- GREENTREE, M. R., LI, Z. X., LI, X. H. & WU, H. 2006. Late Mesoproterozoic to earliest Neoproterozoic basin record of the Sibao orogenesis in western South China and relationship to the assembly of Rodinia. *Precambrian Research* **151**, 79–100.
- GUAN, J. L., ZHENG, L. L., LIU, J. H., SUN, Z. M. & CHENG, W. H. 2011. Zircons SHRIMP U–Pb dating of dolerite from Hekou, Sichuan Province, China and its geological significance. *Acta Geologica Sinica* **85**, 482–90 (in Chinese with English abstract).
- HOFFMAN, P. F. 1989. Speculations on Laurentia's first gigayear (2.0 to 1.0 Ga). *Geology* **17**, 135–8.
- HOFFMAN, P. F. 1997. Tectonic genealogy of North America. In *Earth Structure: An Introduction to Structural Geology and Tectonics* (eds B. A. Van der Pluijm & S. Marshak), pp. 459–64. New York: McGraw-Hill.
- HOU, G., LI, J. H., YANG, M. H., YAO, W. H., WANG, C. C. & WANG, Y. X. 2008a. Geochemical constraints on the tectonic environment of the Late Paleoproterozoic mafic dyke sarms in the North China Craton. *Gondwana Research* **13**, 103–16.
- HOU, G., SANTOSH, M., QIAN, X., LISTER, G. S. & LI, J. 2008b. Configuration of the Late Paleoproterozoic supercontinent Columbia: insights from radiating mafic dyke sarms. *Gondwana Research* **14**, 395–409.
- HUA, R. M. 1990. On the Kunyang Aulacogen. *Acta Geologica Sinica* **64**, 289–301 (in Chinese with English abstract).
- INGLE, S., WEIS, D., SCOATES, J. S. & FREY, F. A. 2002. Relationship between the early Kerguelen plume and continental flood basalts of the paleo-eastern Gondwanan margins. *Earth and Planetary Science Letters* **197**, 35–50.
- LI, Z. X., BOGDANOVA, S. V., COLLINS, A. S., DAVIDSON, A., DE WAELE, B., ERNST, R. E., FITZSIMONS, I. C. W., FUCH, R. A., GLADKUCHUB, D. P., JACOBS, J., KARLSTROM, K. E., LU, S., NATAPOVM, L. M., PEASE, V., PISAREVSKY, S. A., THRANE, K. & VERNIKOVSKY, V. 2008. Assembly, configuration, and break-up history of Rodinia: a synthesis. *Precambrian Research* **160**, 179–210.
- LI, X. H., LI, Z. X., SINCLAIR, J. A., LI, W. X. & CARTER, G. 2006. Revisiting the “Yanbian Terrane”: implications for Neoproterozoic tectonic evolution of the western Yangtze Block, South China. *Precambrian Research* **151**, 14–30.
- LI, X. H., LI, W. X., WANG, X. C., LI, Q. L., LIU, Y., TANG, G. Q., GAO, Y. Y. & WU, F. Y. 2010. SIMS U–Pb zircon geochronology of porphyry Cu–Au–(Mo) deposits in the Yangtze River Metallogenic Belt, eastern China: magmatic response to early Cretaceous lithospheric extension. *Lithos* **119**, 427–38.
- LI, Z. X., LI, X. H., ZHOU, H. & KINNY, P. D. 2002. Grenvillian continental collision in South China: new SHRIMP U–Pb zircon results and implications for the configuration of Rodinia. *Geology* **30**, 163–6.
- LI, X. H., LIU, Y., LI, Q. L. & GUO, C. H., CHAMBERLAIN, K. R. 2009. Precise determination of Phanerozoic zircon Pb/Pb age by multi-collector SIMS without external standardization. *Geochemistry Geophysics Geosystems* **10**, Q04010, doi: [10.1029/2009GC002400](https://doi.org/10.1029/2009GC002400).
- LI, Z. X., WARTHIO, J. A., OCCHIPINTI, S., ZHANG, C. L., LI, X. H., WANG, J. & BAO, C. M. 2007. Early history of the eastern Sibao orogen (South China) during the assembly of Rodinia: new $^{40}\text{Ar}/^{39}\text{Ar}$ dating and U–Pb SHRIMP detrital zircon provenance constraints. *Precambrian Research* **159**, 74–94.
- LI, Z. X., ZHANG, L. & POWELL, C. M. C. A. 1995. South China in Rodinia: part of the missing link between Australia–East Antarctica and Laurentia? *Geology* **23**, 407–10.
- LIGHTFOOT, P. C., HAWKESWORTH, C. J., HERGT, J., NALDRETT, A. J., GORBACHEV, N. S., FEDORENKO, V. A. & DOHERTY, W. 1993. Remobilisation of the continental

- lithosphere by a mantle plume: major-, trace-element, and Sr-, Nd-, and Pb-isotope evidence from picritic and tholeiitic lavas of the Noril'sk District, Siberian Trap, Russia. *Contributions to Mineralogy and Petrology* **114**, 171–88.
- LOUCKS, R. R. 1990. Discrimination of ophiolitic from nonophiolitic ultramafic-mafic allochthons in orogenic belts by the Al/Ti ratio in clinopyroxene. *Geology* **18**, 346–9.
- LUOMAIR, G. W. & HARTI, K. 1978. Lunar initial $^{143}\text{Nd}/^{144}\text{Nd}$: differential evolution of the lunar crust and mantle. *Earth and Planetary Science Letters* **39**, 349–57.
- MEERT, J. G. 2012. What's in a name? The Columbia (Paleopangaea/Columbia) supercontinent. *Gondwana Research* **21**, 987–93.
- MOU, C. L., LIN, S. L. & YU, Q. 2003. The U–Pb ages of the volcanic rock of the Tianbaoshan formation, Huili, Sichuan province. *Journal of Stratigraphy* **27**, 216–9 (in Chinese with English abstract).
- ORMEROD, D. S., HAWKESWORTH, C. J., ROGERS, N. W., LEEMAN, W. P. & MENZIES, M. A. 1988. Tectonic and magmatic transitions in the western Great Basin, USA. *Nature* **333**, 349–53.
- PACES, J. B. & BELL, K. 1989. Non-depleted sub-continental mantle beneath the Superior Province of the Canadian Shield: Nd–Sr isotopic and trace element evidence from midcontinent rift basalts. *Geochimica et Cosmochimica Acta* **53**, 2023–35.
- PENG, M., WU, Y. B., GAO, S., ZHANG, H. F., WANG, J., LIU, X. C., GONG, H. J., ZHOU, L., HU, Z. C., LIU, Y. S. & YAN, H. L. 2012. Geochemistry, zircon U–Pb age and Hf isotope compositions of Paleoproterozoic aluminous A-type granites from the Kongling terrain, Yangtze Block: constraints on petrogenesis and geologic implications. *Gondwana Research* **22**, 140–51.
- PENG, M., WU, Y. B., WANG, J., JIAO, W. F., LIU, X. C. & YANG, S. H. 2009. Paleoproterozoic mafic dyke from Kongling terrain in The Yangtze Craton and its implication. *Chinese Science Bulletin* **54**, 1098–104.
- PENG, P., ZHAI, M.-G., ERNST, R., GUO, J.-H., LIU, F. & HU, B. 2008. A 1.78 Ga Large Igneous Province in the North China craton: the Xiong'er Volcanic Province and the North China dyke swarm. *Lithos* **101**, 260–80.
- PENG, P., ZHAI, M. G. & GUO, J. H. 2006. 1.80–1.75 Ga mafic dyke swarms in the central North China craton: implications for a plume-related break-up event. In *Dyke Swarms—Time Markers of Crustal Evolution* (eds E. Hanski, S. Mertanen, T. Ramö & J. Vuollo). Oxford: Taylor & Francis.
- PENG, P., ZHAI, M. G., GUO, J. H., KUSKY, T. & ZHAO, T. P. 2007. Nature of mantle source contributions and crystal differentiation in the petrogenesis of the 1.78 Ga mafic dykes in the central North China craton. *Gondwana Research* **12**, 29–46.
- PIK, R., DENIEL, C., COULON, C., YIRGU, G., HOFMANN, C. & AYALEW, D. 1998. The northwestern Ethiopian plateau flood basalts: classification and spatial distribution of magma types. *Journal of Volcanology and Geothermal Research* **81**, 91–111.
- POLAT, A., HOFMANN, A. W. & ROSIN, G. M. T. 2002. Boninite-like volcanic rocks in the 3.7–3.8 Ga Isua greenstone belt, West Greenland: geochemical evidence for intraoceanic subduction zone processes in the early Earth. *Chemical Geology* **184**, 231–54.
- QI, L., HU, J. & REGOIRE, D. C. 2000. Determination of trace elements in granites by inductively coupled plasma mass spectrometry. *Talanta* **51**, 507–13.
- ROGERS, J. J. W. & SANTOSH, M. 2002. Configuration of Columbia, a Mesoproterozoic supercontinent. *Gondwana Research* **5**, 5–22.
- RUDNICK, R. L. & FOUNTAIN, D. M. 1995. Nature and composition of the continental crust: a lower crustal perspective. *Reviews of Geophysics* **33**, 267–309.
- SBG (Sichuan Bureau of Geology). 1967. A report of regional geological survey in Huili area of the people's republic of China (the scale of 1:200000) (in Chinese).
- SBGMR (Sichuan Bureau of Geology and Mineral Resources). 1991. *Regional Geology of Sichuan Province*. Beijing: Geology Publishing House, 730 pp (in Chinese with English abstract).
- SUN, M., CHEN, N., ZHAO, G., WILDE, S. A., YE, K., GUO, J., CHEN, Y. & YUAN, C. 2008. U–Pb zircon and Sm–Nd isotopic study of the Huangtuling granulite, Dabie-Sulu belt, China: implication for the Paleoproterozoic tectonic history of the Yangtze Craton. *American Journal of Science* **308**, 469–83.
- SUN, S. S. & McDONOUGH, W. F. 1989. Chemical and isotopic systematics of oceanic basalts: implications for mantle composition and processes. In *Magmatism in the Ocean Basins* (eds A. D. Saunders & M. J. Norry), pp. 313–45. Geological Society of London, Special Publication no. 42.
- SUN, Z. M., YIN, F. G., GUAN, J. L., LIU, J. H., LI, J. M., GENG, Y. R. & WANG, L. Q. 2009. SHRIMP U–Pb dating and its stratigraphic significance of tuff zircons from Heishan formation of Kunyang Group, Dongchuan area, Yunnan Province, China. *Geological Bulletin of China* **28**, 896–900 (in Chinese with English abstract).
- WANG, W. & ZHOU, M. F. 2014. Provenance and tectonic setting of the Paleo- to Mesoproterozoic Dongchuan Group in the southwestern Yangtze Block, South China: implication for the breakup of the supercontinent Columbia. *Tectonophysics* **610**, 110–27.
- WANG, L. J., GRIFFIN, W. L., YU, J. H. & O'REILLY, S. Y. 2010. Precambrian crustal evolution of the Yangtze Block tracked by detrital zircons from Neoproterozoic sedimentary rocks. *Precambrian Research* **177**, 131–44.
- WANG, L. J., YU, J. H., GRIFFIN, W. L. & O'REILLY, S. Y. 2012. Early crustal evolution in the western Yangtze Block: evidence from U–Pb and Lu–Hf isotopes on detrital zircons from sedimentary rocks. *Precambrian Research* **222–223**, 368–85.
- WINGATE, M. T. D. & COMPSTON, W. 2000. Crystal orientation effects during ion microprobe U–Pb analysis of baddeleyite. *Chemical Geology* **168**, 75–97.
- WOOD, D. A., JORON, J. L. & TREUIL, M. 1979. A re-appraisal of the use of trace elements to classify and discriminate between magma series erupted in different tectonic settings. *Earth and Planetary Science Letters* **45**, 326–36.
- WU, M. D., DUAN, J. S., SONG, X. L., CHEN, L. & DAN, Y. 1990. *Geology of Kunyang Group in Yunnan Province*. Kunming: Scientific Press of Yunnan Province, 265 pp (in Chinese with English abstract).
- WU, Y. B., ZHENG, Y. F., GAO, S., JIAO, W. F. & LIU, Y. S. 2008. Zircon U–Pb age and trace element evidence for Paleoproterozoic granulite-facies metamorphism and Archean crustal rocks in the Dabie Orogen. *Lithos* **101**, 308–22.
- XIONG, Q., ZHENG, J. P., YU, C. M., SU, Y. P., TAN, G. H. Y. & ZHANG, Z. H. 2009. Zircon U–Pb age and Hf isotope of Quanyishang A-type granite in Yichang: significance for the Yangtze continental cratonization in Paleoproterozoic. *Chinese Science Bulletin* **54**, 436–46.
- XU, Y. G., CHUNG, S. L., JAHN, B. M. & WU, G. Y. 2001. Petrologic and geochemical constraints on the

- petrogenesis of Permian–Triassic Emeishan flood basalts in southwestern China. *Lithos* **58**, 145–68.
- YAO, J., SHU, L. & SANTOSH, M. 2011. Detrital zircon U–Pb geochronology, Hf-isotopes and geochemistry — new clues for the Precambrian crustal evolution of Cathaysia Block, South China. *Gondwana Research* **20**, 553–67.
- YIN, F. G., SUN, Z. M. & ZHANG, Z. 2011. Mesoproterozoic stratigraphic-structure framework in Huili-Dongchuan area. *Geology Review* **57**, 770–8 (in Chinese with English abstract).
- ZHANG, C. H., GAO, L. Z., WU, Z. J., SHI, X. Y., YAN, Q. R. & LI, D. J. 2007. SHRIMP U–Pb zircon age of tuff from the Kunyang group in central Yunnan: evidence for Grenvillian orogeny in south China. *Chinese Science Bulletin* **52**, 1517–25.
- ZHANG, S., LI, Z. X., EVANS, D. A. D., WU, H. & LI, H. 2012. Pre-Rodinia supercontinent Nuna shaping up: a global synthesis with new paleomagnetic results from North China. *Earth and Planetary Science Letters* **353–354**, 145–55.
- ZHANG, S. B., ZHENG, Y. F., WU, Y. B., ZHAO, Z. F., GAO, S. & WU, F. Y. 2006a. Zircon isotope evidence for ≥ 3.5 Ga continental crust in the Yangtze craton of China. *Precambrian Research* **146**, 16–34.
- ZHANG, S. B., ZHENG, Y. F., WU, Y. B., ZHAO, Z. F., GAO, S. & WU, F. Y. 2006b. Zircon U–Pb age and Hf–O isotope evidence for Paleoproterozoic metamorphic event in South China. *Precambrian Research* **151**, 265–88.
- ZHAO, G. C., CAWOOD, P. A., WILDE, S. A. & SUN, M. 2002a. Review of global 2.1–1.8 Ga orogens: implications for a pre-Rodinia supercontinent. *Earth-Science Reviews* **59**, 125–62.
- ZHAO, G. C., SUN, M., WILDE, S. A. & LI, S. Z. 2004. A Paleo-Mesoproterozoic supercontinent: assembly, growth and breakup. *Earth-Science Reviews* **67**, 91–123.
- ZHAO, X. F. & ZHOU, M. F. 2011. Fe–Cu deposits in the Kangdian region, SW China: a Proterozoic IOC G(iron–oxide–copper–gold) metallogenic province. *Mineralium Deposita* **46**, 731–47.
- ZHAO, X. F., ZHOU, M. F., LI, J. W., SUN, M., GAO, J. F., SUN, W. H. & YAN, J. H. 2010. Late Paleoproterozoic to early Mesoproterozoic Dongchuan Group in Yunnan, SW China: implications for tectonic evolution of the Yangtze Block. *Precambrian Research* **182**, 57–69.
- ZHAO, X. F., ZHOU, M. F., LI, J. W. & QI, L. 2013. Late Paleoproterozoic sedimentary rock-hosted stratiform copper deposits in South China: their possible link to the supercontinent cycle. *Mineralium Deposita* **48**, 129–36.
- ZHAO, T. P., ZHOU, M. F., ZHAI, M. G. & XIA, B. 2002b. Paleoproterozoic rift-related volcanism of the Xiong'er group, North China craton: implications for the breakup of Columbia. *International Geology Review* **44**, 336–51.
- ZHENG, J. P., GRIFFIN, W. L., O'REILLY, S. Y., ZHANG, M., PEARSON, N. & PAN, Y. 2006. Widespread Archean basement beneath the Yangtze Craton. *Geology* **34**, 417–20.
- ZHOU, M. F., ZHAO, X. F., CHEN, W., LI